

Prevalence of Cardiovascular Disease Risk Factors among Adults in Slum of Dhaka City

Dr. S. M. Zahid Hassan Arefin

Master of Philosophy (M.Phil)

Epidemiology & Biostatistics

Session: 2011-2012

Registration No: 293

498070



BIHS

**Department of Epidemiology & Biostatistics
Bangladesh Institute of Health Sciences (BIHS)
Faculty of Biological Science
University of Dhaka**

R
616.1
ARP

file

070022

DECLARATION

I, hereby humbly declare that the thesis work entitled “**Prevalence of Cardiovascular Disease Risk Factors among Adults in Slum of Dhaka City**”, a requirement for the degree of Master of Philosophy (M.Phil) in Epidemiology & Biostatistics from Bangladesh Institute of Health Sciences (BIHS), under faculty of Biological Science, University of Dhaka, was carried out by me under the guidance of Dr Saidur Rahman Mashreky, Associate Professor, Department of Epidemiology and Biostatistics, Bangladesh Institute of Health Sciences (BIHS) during the period of June 2014 to July 2015.

No part of the work has been submitted elsewhere for any other purpose.



.....
S. M. Zahid Hassan Arefin

M.Phil in Epidemiology & Biostatistics

Session: 2011- 2012

Bangladesh Institute of Health Sciences (BIHS)

Dhaka, Bangladesh.


498070

CERTIFICATE

This is to certify that the thesis entitled “**Prevalence of Cardiovascular Disease Risk Factors among Adults in Slum of Dhaka City**”, submitted by S. M. Zahid Hassan Arefin in partial fulfillment of the requirement for the degree of Master of Philosophy (M.Phil) in Epidemiology & Biostatistics from Bangladesh Institute of Health Sciences (BIHS), University of Dhaka, was carried out under my supervision.

To the best of my knowledge no part of the work has been submitted for another degree or qualification in any other institute.

Supervisor



.....
Dr Saidur Rahman Mashreky. MPH, Dip. MEd., PhD.


Associate Professor

Dept. of Epidemiology and Biostatistics

Bangladesh Institute of Health Sciences (BIHS),

University of Dhaka.

.. 498070



University of Dhaka
Faculty of Biological Science

The undersigned certify that they have carefully read and examined the student on this thesis and being satisfied, recommended the faculty of Biological Science, University of Dhaka, for acceptance of this thesis entitled "**Prevalence of Cardiovascular Disease Risk Factors among Adults in Slum of Dhaka City**", by S. M. Zahid Hassan Arefin in partial fulfillment of the requirement for the degree of Master of Philosophy (M.Phil) in Epidemiology & Biostatistics for the session 2011-2012.

Board of Examiners

Chairman

Signature.....

Name:

Designation:

Member


498070

Signature.....

Name:

Designation:

Member



Signature.....

Name:

Designation:

Member

Signature.....

Name:

Designation:

Date of approval:

Dedicated to

*My father late M.A. Quashem, Mother Monwara
Begum, My wife Dr. Farhana Jahan, My kids Aruba &
Aariz*

&

*My respected Teacher
Dr Saidur Rahman Mashreky*

ACKNOWLEDGEMENTS

First of all, I want to express my gratitude to the Almighty Allah, the most merciful for giving me the passion to complete this study.

I would like to take the opportunity to pay my respect and heartfelt gratitude to my supervisor, Dr Saidur Rahman Mashreky, Associate Professor, Dept. of Epidemiology, Bangladesh Institute of Health Sciences (BIHS) for his excellent guidance, directions, valuable suggestions, time and intellectual ideas throughout the study that always inspired me to take the challenges with confidence.

I find my words inadequate to express my heartfelt gratitude to my co-guide Shirin Jahan Mumu, Assistant Professor, Dept. of Epidemiology, Bangladesh Institute of Health Sciences (BIHS) for her continuous help, support and valuable suggestions in each and every stage of this research.

I wish to express my sincere thanks to Professor M.A. Hafez, Head, Department of Biostatistics, BIHS for giving me his valuable time, suggestion and intellectual inputs.

I also wish to thank specially to Professor Abidul Haque, Head, Department of Epidemiology and Biostatistics, Institute of Child and Mother Health for his continuous, invaluable support.

In addition, I want to give my special, heartfelt gratitude and thanks to Dr. Morshed Alam and Dr. Shahebzada for their great, laborious effort in helping me to collect data by going from door to door in the slum during the hottest time of year.

Finally I am extremely grateful to my wife for her unflagging support and sacrifice throughout the study and helping me in every possible way to the completion of this work.

Last of all, I am grateful to all my participants who were kind enough to spend their time in providing data for this study.

Dr. S.M Zahid Hassan Arefin

CONTENTS

LIST OF CONTENTS	PAGE NO
List of Tables	iii
List of abbreviations	iv
Abstract	v- i
CHAPTER 1	
INTRODUCTION	1-11
1.1 Background	1-8
1.2 Justification of the study	9
1.3 Research question	10
1.4 Objectives of the study	11
CHAPTER 2	
LITERATURE REVIEW	12-25
2.1 Global cardiovascular mortality	12
2.2 Cardiovascular disease in India	13-14
2.3 Cardiovascular disease in Bangladesh	14
2.4 Coronary heart disease risk of slum dwelling residents	14
2.5 Individual risks for CVD	15-16
2.6 Major proximal risk factors for CVD	16-25
CHAPTER 3	
SUBJECTS AND METHODS	26- 38
3.1 Study design	26
3.2 Study period	26
3.3 Study area	26

3.4	Study population	26
3.5	Study sample	26
3.6	Inclusion criteria	26
3.7	Exclusion criteria	27
3.8	Sample size	27
3.9	Sampling technique	27
3.10	Data collection techniques/methods	28
3.11	Data collection tools	29
3.12	Data analysis	30
3.13	List of variables	31
3.14	Conceptual Framework	32
3.15	Operational Definitions	33-38
3.16	Ethical Consideration	38
CHAPTER 4		
RESULTS		39-58
CHAPTER 5		
DISCUSSION		59-63
CHAPTER 6		
CONCLUSION, LIMITATION AND RECOMMENDATIONS OF THE STUDY		64-66
6.1	Conclusion	64
6.2	Limitation	65
6.3	Recommendation	66
CHAPTER 7		
REFERENCES		67-77

CHAPTER 8

ANNEXURE

	78-108
Annex 1 Work plan	78
Annex 2 Informed Consent Form	79-80
Annex 3 Questionnaire	81-104
Annex 4 Show cards	105-108

List of Tables

Table No	Title	Page No
Table 4.1.1	Socio-demographic characteristics of study subjects- age, marital status, education, occupation, economic status, religion, family size & house type	40
Table 4.1.2	Behavioral characteristics of the respondents	42
Table 4.1.3	Dietary pattern- fruit, vegetable, oil and salt consumption of the respondents	45
Table 4.1.4	Biological risk factors among the respondents	47
Table 4.1.5	Family history of coronary heart disease/heart attack and stroke	49
Table 4.2.1	Association between hypertension and socio-demographic characteristics	51
Table 4.2.2	Association between hypertension and modifiable risk factors	52
Table 4.2.3	Association between hypertension and family history of heart attack or stroke	53
Table 4.3.1	Effect of various modifiable and non-modifiable risk factors on hypertension	55
Table 4.4.1	Effect of sex and age on current smoking status	56
Table 4.4.2	Effect of sex and age on current smokeless tobacco user	56
Table 4.4.3	Effect of sex and age on physical activity	57
Table 4.4.4	Effect of sex and age on overweight and obesity	57
Table 4.4.5	Effect of sex and age on waist circumference	58
Table 4.4.6	Effect of sex and age on diabetes	58

List of Abbreviations

AIDS	Acquired Immuno Deficiency Syndrome
BIHS	Bangladesh Institute of Child Health
BMI	Body Mass Index
BP	Blood Pressure
CHD	Coronary Heart Disease
CI	Confidence Interval
cm	Centimeter
CVD	Cardiovascular disease
DALY	Disability adjusted life year
DBP	Diastolic Blood Pressure
DM	Diabetes Mellitus
gm	Gram
GPAQ	Global Physical Activity Questionnaire
HDL	High Density Lipoprotein
HIV	Human Immuno deficiency Virus
HTN	Hypertension
IHD	Ischemic Heart Disease
Kg	Kilogram
LDL	Low Density Lipoprotein
MET	Metabolic Equivalent
MI	Myocardial Infarction
ml	Mililiter
mmol/L	Milimole per liter
NCD	Non-Communicable Diseases
OR	Odds Ratio
SBP	Systolic Blood Pressure
SD	Standard Deviation
SPSS	Statistical Package for Social Sciences
WC	Waist Circumference
WHO	World Health Organization

Abstract

Background: Cardiovascular disease (CVD) is one of the leading causes of death worldwide and hypertension is the leading modifiable risk factor for cardiovascular mortality. Slum living communities are increasingly adopting urban lifestyles and becoming more vulnerable to risk of developing CVD. Few studies have so far focused on this aspect. **Objective:** The aim of this study was to estimate the prevalence of risk factors of cardiovascular disease among adults in a slum of Dhaka city. **Methods:** It was a cross sectional survey conducted on conveniently selected 461 people at Agargaon slum area and this area was selected purposively. Face to face interview was taken by using structured questionnaire. **Results:** Total 461 adult slum people were studied, of them males were 241(52.3%) and females were 220(47.7%). Mean age of the respondents was 43.04 (SD±8.86) years. More than half of study subjects (57%) had no formal education. Majority of the females (81.8%) were house hold worker. About one third (32.8%) males were small businessmen; day laborer was about 12%. Driver and rickshaw pullers were all males, 12% and 7.5% respectively. The mean monthly family income of the respondents was 13754 (SD±4592) taka. Overall prevalence of smoking tobacco and smokeless tobacco use was 39.3% and 47.5% respectively. Based on MET minutes/week, 36.2% of the study subjects fell into low physical activity category. Most of the METs of women were contributed by moderate activities. Almost ninety per cent study subjects (89.6%) were lifetime abstainer of alcohol. None of the study subjects consumed adequate fruit or vegetables on an average day. Nine percent male consumed more than recommended level of oil daily, whereas it was 35.2% in case of female. Mean salt consumption per day was 13.2 (SD± 3.79) grams, which was more than double of the recommended level and 99% participants consume salt in excess of their normal requirement. Mean BMI of respondents was 22.37 (SD±2.92). Prevalence of overweight and obesity was 15.6%. Mean waist circumferences was 79.83 (SD±7.71) cm. The prevalence of high waist circumference was significantly higher in females than in males (45.5% versus 12.4%). Overall prevalence of hypertension was 14.8% and the rates among male and female were 16.6% and 12.7%

respectively. The mean systolic blood pressure of male and female was 117.43 (SD±14.69) mmHg and 112.52 (SD±13.28) mmHg. A little less than one third (26.7%) of the study subjects never measured their blood pressure. Among the currently diagnosed hypertensive, 64.7% were unaware of their high blood pressure. More than one third (37.5%) of subjects who were on antihypertensive treatment, had blood pressure within normal level. Around 87.6% of participants never measured their blood glucose and about 5% of the people were previously diagnosed to have diabetes. Statistical significant association was found between hypertension and advancing age, higher monthly family income, increased BMI, low physical activity, smokeless tobacco use, diabetes and increased waist circumference in male sex. Logistic regression showed increased risk of hypertension among male sex, higher age group, high income group, smokeless tobacco use, low physical activity, high BMI and high waist circumference. **Conclusion:** The study population showed high prevalence of hypertension and other cardiovascular disease risk factors like both smoking and smokeless tobacco use, low physical activity with very minimum exercise and a considerable amount of time of sedentary activity, very minimum fruit and vegetable intake with very high salt intake. Majority of hypertensive patients were unaware about their hypertension status. There is need for greater awareness of hypertension and other cardiovascular disease risk factors in this population.

CHAPTER - 1
INTRODUCTION

Chapter 1

Introduction

1.1 Background

Cardiovascular disease (CVD) is a major health problem throughout the world and a common cause of premature morbidity and mortality. According to World Health Organization (WHO), CVDs are the number one cause of death globally. An estimated 17.5 million people died from CVDs in 2012, representing 31% of all global deaths (1). Of these deaths, an estimated 7.4 million were due to coronary heart disease and 6.7 million were due to stroke (1). The number of people who die from CVDs, mainly from heart disease and stroke, will increase to reach 23.3 million by 2030 (2,3). CVDs are projected to remain the single leading cause of death (2). Over 80% of CVD deaths take place in low- and middle-income countries (3). It was found as the leading cause of death in population older than 45 years in sub-Saharan Africa (4). South Asians represent one of the largest and fastest growing ethnic groups in the world and they are experiencing the epidemics of cardiovascular disease. In Bangladesh, CVD accounts for 27% of deaths due to all causes, whereas non-communicable diseases (NCD) are estimated to account for 52% of all deaths (5). Bangladesh national survey on NCD risk factors 2010 showed 17.6% people (both sexes) overweight (BMI) ≥ 25 kg/m², 21.7% had increased waist circumference (men ≥ 94 cm, women ≥ 80 cm, prevalence of hypertension 17.9% and diabetes 3.9% (6). A systematic review and meta-analysis in Bangladesh showed the pooled hypertension and type 2 diabetes mellitus prevalence were 13.7% (12.1%–15.3%) and 6.7% (4.9%–8.6%), respectively (7). Another hospital based study in Bangladesh done among patients with probable cardiovascular disease showed lipid level disorder (47.05%), hypertension (28.05%), heart failure (27.25%), ischemic heart disease (21.55%) and 40.39% were associated with diabetes (8).

Cardiovascular disease (CVD) accounts for half of non-communicable disease deaths worldwide (9). Although CVD risk is perceived to be low in economically developing

countries, ischemic heart disease and stroke were 2 of the 3 leading causes of mortality in such countries in 1990 (10). In China, CVD accounted for nearly 40% of all deaths in 1994 (11). Furthermore, CVD incidence and mortality in China are projected to increase substantially during the next 20 years. Although the prevalence of some CVD risk factors has decreased in economically developed countries, the corresponding prevalence has increased in economically developing countries (11). For example, in China, the prevalence of hypertension increased from 11.3% in 1991 to 27.2% in 2000 (12,13). Furthermore, mean cholesterol levels have increased during the past decade, with the most recent report indicating that 23.8% and 9.0% of the Chinese adult population have a total cholesterol level between 5.2 mmol/L and 6.2 mmol/L and ≥ 6.2 mmol/L, respectively (14). China is currently undergoing rapid demographic, social, and economic changes that may further increase the burden of CVD.

What is CVD

CVD is a general category of diseases that affects the heart and the circulatory system. CVD is caused by disorders of the heart and blood vessels, and includes coronary heart disease (CHD), congestive heart failure (CHF), stroke, hypertension, peripheral artery disease, and rheumatic heart disease, congenital heart disease, deep vein thrombosis and pulmonary embolism (1,8). CHD alludes to a reduction of blood flow due to thickening and hardening of the arteries that supply the heart muscle. A complete cut off of the blood supply results in the death of heart cells, and a heart attack (MI) occurs. CHF is a disorder where the heart loses its ability to pump blood efficiently. Finally stroke occurs when a blood vessel bringing oxygen and nutrients to the brain bursts or is clogged by a blood clot or some other particle (15).

Some well documented cardiovascular disease risk factors are as follows:

A. Major modifiable risk factors

1. Major behavioral risk factors (16)

The major behavioral risk factors identified in the World Health Report 2002 are:

- Tobacco use
- Unhealthy diet (low fruit and vegetable; high salt and oil consumption)
- Physical inactivity
- Harmful alcohol consumption

2. Major biological risk factors (16)

The major biological risk factors identified in the World Health Report 2002 are:

- Overweight and obesity
- Raised blood pressure
- Raised blood glucose
- Abnormal blood lipids and its subset raised total cholesterol

Another important modifiable risk factor not falling into these two categories-

- low socioeconomic status.

B. Non-modifiable risk factors

- Advancing age
- Heredity or family history
- Gender
- Ethnicity or race

Major behavioral risk factors:

1. Tobacco use

About 1.3 billion people worldwide smoke and the number of smokers continue to rise (16). Among these, about 84% live in developing and transitional economy countries (16,17). Tobacco is the fourth most common risk factor for disease and the second major cause of death worldwide. It is currently responsible for the death of one in ten adults worldwide (about 4.9 million deaths each year) (16). In Bangladesh, overall prevalence of smoking is 26.2% (54.8% in men and 1.3% in women) (6). Overall, smokers have a 70% greater level of CVD risk than nonsmokers; persons who smoke greater than or equal to 2 packs of cigarettes per day have a two- to threefold greater risk for CVD (18). The risk for CVD also increases with greater depth of inhalation and with increasing years of smoking, although persons who stop smoking eventually reduce their risk for CVD to a level approaching that of nonsmokers (18). Cigarette smoking has been reported to act synergistically with other known risk factors for CVD.

2. Unhealthy diet - low fruit and vegetable, high salt and oil consumption

Low intake of fruits and vegetables is estimated to cause about 31% of ischemic heart disease and 11% of stroke worldwide (19). Overall, 2.7 million lives could potentially be saved each year worldwide if fruit and vegetable consumption were increased (20). Of the burden attributable to low fruit and vegetable intake, about 85% was from cardiovascular diseases (19). Adequate consumption of fruit and vegetables reduces the risk for cardiovascular diseases. In our country, 95.7% people do not consume adequate fruit or vegetables on an average day (6). High dietary intakes of saturated fat, trans-fats and salt are linked to cardiovascular risk (21). The amount of dietary salt consumed is an important determinant of blood pressure levels and overall cardiovascular risk. High consumption of saturated fats and trans-fatty acids is linked to heart disease; elimination of trans-fat and replacement of saturated with polyunsaturated vegetable oils lowers coronary heart disease risk (21).

3. Physical Inactivity

Physical inactivity causes about 1.9 million avoidable deaths per year worldwide (19). Physically inactive persons have a 20% to 30% increased risk of all-cause mortality as compared to those who adhere to 30 minutes of moderate intensity physical activity on most days of the week (16). Globally, physical inactivity accounts for 21.5% of ischemic heart disease and 11% of ischemic stroke (16). According to NCD risk factor survey Bangladesh 2010, 27% of Bangladeshi people fall into low physical activity category (6). A review of 43 epidemiologic studies in 1987 indicated that physical activity reduces the risk of CHD (22). The relative risk for CHD associated with physical inactivity is approximately 1.9, slightly lower than the relative risks associated with increased systolic blood pressure (2.1), cigarette smoking (2.5), and elevated serum cholesterol levels (2.4) (18). Several studies indicate that endurance exercise training among patients with documented CHD is associated with reduced morbidity and mortality and that physical activity might improve the likelihood of survival from a myocardial infarction (i.e. heart attack) (23). In addition, evidence documents an association between regular, moderate-intensity physical activity and the lowering of several other risk factors for CVD, including blood lipid levels, resting blood pressure among persons with borderline hypertension, body composition and overweight, and glucose tolerance and insulin sensitivity (24).

4. Harmful alcohol consumption

In 2000, alcohol use caused 3.2% of deaths (1.8 million) worldwide, and 4% of the global disease burden (19). Alcohol consumption is the leading risk factor for disease burden in developing countries and the third largest risk factor in developed countries (25). Heavy alcohol use increases the risk of cardiovascular disease and stroke (16). Though people of our country is very much religious and ninety four per cent adult in Bangladesh are lifetime abstainer of alcohol, the rates of alcohol and drug abuse is relatively high in slums of Dhaka (6,26).

Major biological risk factors:

1. Overweight and obesity

At least 2.6 million people die each year as a result of being overweight or obese (20). The high death rate might occur largely as a consequence of the influence of overweight on blood pressure, blood lipid levels, and the onset of diabetes (27). Risks of coronary heart disease, ischemic stroke and type 2 diabetes mellitus increase steadily with increasing BMI (19). A report from the Framingham Study indicates that overweight is also an independent risk factor for CVD (18). With rare exceptions, overweight develops from eating too much and exercising too little. The prevalence of overweight has increased substantially in the U.S. population during the last 10 years (18). In Bangladesh, proportion of overweight in women (21.6%) exceeded the proportion of those in men (13%) (6). Waist circumference is an approximate index of intra-abdominal fat mass and total body fat. In our country, eight per cent men and 33.7% women (21.7% sexes combined) had increased waist circumference (6). Changes in waist circumference reflect changes in risk factors for cardiovascular disease.

2. Hypertension (Raised blood pressure)

Worldwide, raised blood pressure is estimated to cause 7.1 million deaths, about 13% of the total (16). High blood pressure is a major risk factor for CVD. Some evidence documents that blood pressure-related risk for CVD increases continuously from lowest to highest values for either systolic or diastolic blood pressure (18). Elevated blood pressure is often associated with other well-known risk factors, including dietary intake, elevated blood lipid levels, obesity, smoking, diabetes mellitus, and physical inactivity (18). The prevalence of documented hypertension in Bangladesh was 12.5% (men 10.9% and women 13.9%) (6).

3. Raised blood glucose

It is predicted that the number of people living with diabetes is expected to rise from 366 million in 2011 to 552 million by 2030, if no urgent action is taken (28). The risk for CVD is three times as high among diabetic women as it is among women without

diabetes mellitus. Similarly, the risk for CVD is twice as high among diabetic men as it is among men without diabetes mellitus (18,29). Impaired glucose tolerance and impaired fasting glycaemia, also called pre-diabetes condition are risk categories for future development of diabetes and cardiovascular disease (30). The age-adjusted mortality, mostly due to coronary heart disease in many populations, is 2-4 times higher in diabetic than in the non-diabetic population (16). People with diabetes have a twofold increase risk of stroke (16). In Bangladesh, the age-adjusted prevalence of diabetes and pre-diabetes is quite high, 9.7% and 22.4%, respectively, and there is no significant difference between the sexes (31).

4. Abnormal blood lipids

Raised cholesterol is estimated to cause 18% of the global cerebrovascular disease and 56% of global ischemic heart disease (19). Levels of plasma HDL cholesterol are inversely related to coronary artery disease incidence, and the relationship is independent of total cholesterol, LDL and triglyceride levels (32). Increased triglycerides is an independent risk factor for coronary heart disease after controlling for LDL and HDL cholesterol (33). No study has so far been done in our country regarding the lipid status of our population.

Other modifiable risk factor

1. Low socioeconomic status

There is consistent inverse relationship of lower socioeconomic status with risk of heart disease and stroke (34).

Non-modifiable risk factors

1. Advancing age

Most powerful independent risk factor for cardiovascular disease is older age. Risk of stroke doubles every decade after age 55 (35).

2. Heredity or family history

Increased risk if a first-degree blood relative has had coronary heart disease or stroke before the age of 55 years (for a male relative) or 65 years (for a female relative) (35).

3. Gender

Higher rates of coronary heart disease among men compared with women (premenopausal age); Once past the menopause, a woman's risk is similar to a man's. Risk of stroke is similar for men and women (35).

4. Ethnicity or race

Increased stroke noted for Blacks, some Hispanic Americans, Chinese, and Japanese populations. Increased cardiovascular disease deaths noted for South Asians and American Blacks in comparison with Whites (35).

1.2 Justification

Cardiovascular disease is increasing worldwide. People in low- and middle-income countries are more exposed to risk factors such as less awareness about food and exercise. Prevention effort in low income countries is very few compared to people in high-income countries. People in low- and middle-income countries who suffer from CVDs have less access to effective and equitable health care services. As a result, many people in these countries die from CVDs and other non-communicable diseases, often in their most productive years. At macro-economic level, CVDs place a heavy burden on the economies of low- and middle-income countries. In 2010, the global direct and indirect cost of CVD was approximately US\$ 863 billion and is estimated to rise 22% to US\$ 1,044 billion by 2030. Overall, the cost for CVD alone could be as high as US\$ 20 trillion over the next 20 year period (36). In recent years, several studies in Bangladesh have shown high prevalence of cardiovascular disease and its negative impact on survival but few studies have focused on slum populations. Bangladesh is one of the densely populated countries in the world and Dhaka city's population is about 17.6 million, of which an estimated 3.4 million people live in some 5000 slums (37). These slum population, though they represent a large part of urban population are always neglected. They do not get proper health care and education and they are not even aware of their risk of developing diseases. In this population-based study on persons living in a slum of Dhaka city, we seek to determine the prevalence of risk factors related to cardiovascular disease among slum population, which if properly addressed will help policy makers and health care providers to take appropriate decision regarding this matter. This will also help in future studies on relevant issues.

1.3 Research Question

What are the prevalence of cardiovascular disease risk factors such as tobacco use, unhealthy diet, low physical activity, harmful alcohol consumption, overweight- obesity, hypertension and diabetes among adults in a slum of Dhaka city?

1.4 Objectives

1.4.1 General Objective

- To estimate the prevalence of cardiovascular disease risk factors among adults in a slum of Dhaka city.

1.4.2 Specific Objectives

- To estimate the prevalence of modifiable risk factors of cardiovascular disease such as tobacco use, low fruit-vegetable and high salt-oil consumption, low physical activity, harmful alcohol consumption, overweight-obesity, hypertension, diabetes and low socioeconomic status as reflected by low monthly family income among adults in a slum of Dhaka city.
- To estimate the prevalence of non modifiable risk factors of cardiovascular disease such as advancing age, male sex and family history of heart attack or stroke among adults in a slum of Dhaka city.
- To see the association between hypertension and various socio-demographic characteristics, modifiable and non-modifiable risk factors.
- To find out the effect of various modifiable and non-modifiable risk factors on hypertension.
- To find out the effect of non-modifiable risk factors on various modifiable risk factors.

CHAPTER - 2
LITERATURE REVIEW

Chapter 2

Literature Review

In recent years, the dominance of chronic diseases as major contributors to total global mortality has emerged. By 2005, the total number of cardiovascular disease (CVD) deaths (mainly coronary heart disease, stroke, and rheumatic heart disease) had increased globally to 17.5 million from 14.4 million in 1990 (1). Of these, 7.6 million were attributed to coronary heart disease and 5.7 million to stroke (1). More than 80 percent of the deaths occurred in low and middle income countries (38). The World Health Organization (WHO) estimates there will be about 20 million CVD deaths in 2015, accounting for 30 percent of all deaths worldwide (39). By 2030, researchers project that non-communicable diseases will account for more than three-quarters of deaths worldwide; CVD alone will be responsible for more deaths in low income countries than infectious diseases (including HIV/AIDS, tuberculosis, and malaria), maternal and perinatal conditions, and nutritional disorders combined (40). Thus, CVD is today the largest single contributor to global mortality and will continue to dominate mortality trends in the future (38).

2.1 Global cardiovascular mortality

Globally, there is an uneven distribution of age-adjusted CVD mortality. The lowest age-adjusted mortality rates are in the advanced industrialized countries and parts of Latin America, whereas the highest rates today are found in Eastern Europe and a number of low and middle income countries. Age-standardized mortality rates for CVD are in excess of 500 per 100,000 in Russia and Egypt; between 400 and 450 for South Africa, India and Saudi Arabia; and around 300 for Brazil and China. This is in contrast to rates of between 100 and 200 per 100,000 for Australia, Japan, France, and the United States. Overall, age-adjusted CVD death rates are today higher in major low and middle income countries than in developed countries (41). Examination of coronary heart disease (CHD) mortality trends across countries reveals considerable variability in the shape and

magnitude of CHD epidemics since the 1950s. Trends are not consistent even among countries within the same geographic region. In general, three trending patterns of CHD mortality can be observed: a rise-and-fall pattern where mortality rates increased, peaked, and then fell significantly; a rising pattern, where rates have steadily increased indicating an ongoing epidemic; and a flat pattern, where CHD mortality rates have remained relatively low and stable. The rise-and-fall pattern is most notable in high income Anglo-Celtic, Nordic, and Northwestern Continental European countries as well as in the United States and Australia. In these countries, CHD mortality rates peaked in the 1960s or early 1970s and have since fallen precipitously, by an average of about 50 percent (42). The rising pattern of CHD is most notable in Eastern European and former Soviet countries, where mortality rates have continued to increase at an alarming pace and where the highest mortality rates ever recorded are currently being observed. By contrast, CHD mortality rates in Japan and several European Mediterranean countries have remained relatively low, following the flat pattern (43).

2.2 Cardiovascular disease in India

In developed nations the rise in the burden of CVD occurred over several decades due to a long period of epidemiological transition. In India, perhaps because of the rapid pace of economic development, epidemiological changes have spanned a much shorter time. As a consequence, cardiovascular disease (CVD) has emerged as the leading cause of death all over India, with coronary heart disease (CHD) affecting Indians at least 5-6 years earlier than their western counterparts (44). Current estimates from disparate cross-sectional studies indicate the prevalence of CHD to be between 7-13 per cent in urban and 2-7 per cent in rural India (45). In addition, migration and urbanization have resulted in an increase in the prevalence of risk factors such as diabetes, overweight (46). In India in the past five decades, rates of coronary disease among urban populations have risen from 4% to 11% (47). In India, the leading cause of death is cardiovascular disease. At the same time, it has been found that cardiovascular disease is third overall in the burden of disease, the other two being infectious and parasitic diseases and unintentional injuries. In India, deaths from coronary heart disease rose from 1.17 million in 1990 to 1.59 million in 2000. The prevalence varies by site, age group studied, and diagnostic criteria used,

but an urban prevalence of about 10% in adults aged ≥ 35 years old is a credible estimate based on several surveys (48).

2.3 Cardiovascular disease in Bangladesh

Cardiovascular diseases are one of the major health problems throughout the world. It is emerging as a serious health problem in Bangladesh and other developing countries. Amongst the heart diseases hypertension, rheumatic fever, rheumatic heart diseases, ischemic heart diseases and congenital heart diseases are common. There is a common belief that heart disease is a disease of rich people, which is not correct. Rheumatic fever and rheumatic heart diseases commonly affects poor people living in overcrowding and poverty. Congenital heart diseases and hypertension can affect both rich and poor people. Poor people are not immune from Ischemic heart disease. Latest survey on cardiovascular diseases carried out in Bangladesh showed prevalence of hypertension in adult population about 20-25% and ischemic heart disease about 10% (49). In Bangladesh, CVD accounts for 27% of deaths due to all causes (5). Bangladesh national survey on NCD risk factors 2010 showed prevalence of hypertension 17.9% and (6). A systematic review and meta-analysis in Bangladesh showed that pooled hypertension prevalence were 13.7% (12.1%–15.3%) (7).

2.4 Coronary heart disease risk of slum dwelling residents

Coronary heart disease (CHD) is the top cause of mortality and morbidity in India. People in slums are generally at a higher risk for CHD than Indians living in more affluent areas mostly because of the higher prevalence of major CHD risk factors such as uncontrolled hypertension and tobacco use amongst them. Knowing their CHD risk perceptions and bringing them into line with the actual CHD risk is a prerequisite for effective CHD risk management. Consequently, there is need to develop tailored interventions focusing medication management and tobacco cessation to reduce growing CHD epidemic among slum dwellers and long-term CHD burden in India (50).

2.5 Individual risks for CVD

Proximal risks for CVD include those associated with consumption patterns (mainly linked to diets, tobacco and alcohol use), activity patterns, and health service use as well as biological risk factors such as increased cholesterol, blood pressure, blood glucose, and clinical disease. The Framingham Study first centered attention on the concept of “risk factors” associated with CVD, and most recently reported substantial 30-year risk data showing the accumulation of risk over time (51). Importantly, risk factors for the incidence of CVD and those associated with CVD severity or mortality are not synonymous. Risk factors for incidence become important starting very early in life and accumulate with behavioral, social, and economic factors over the life course to culminate in biological risks for CVD such as increased cholesterol, blood pressure, blood glucose, and clinical disease. Over the past few decades, the effectiveness of early screening and long-term treatment for biological risks or early disease has contributed to the sharp declines in CVD mortality seen in many countries (52).

Relatively few major behavioral and biological risk factors account for CVD incidence around the world. Tobacco use, diet (including alcohol, total calorie intake, and specific nutrients) and physical inactivity serve as the three major behavioral risks. Between them, they account for a significant proportion of cancer, diabetes, and chronic respiratory disease incidence in addition to CVD (53). Concerted action focused on these behavioral risks, along with biological risks such as high blood pressure, high blood lipids, and high blood glucose, would have a wide impact on the global incidence and burden of disease. High blood pressure, tobacco use, elevated blood glucose, physical inactivity, and overweight and obesity are the five leading factors globally. In middle income countries, alcohol replaces high blood glucose in the top five; in low income countries, a lack of safe water, unsafe sex, and under-nutrition are important.

The Global Burden of Disease and Risk Factors report provides additional analysis of the relative contribution of individual risk factors specifically to CVD burden. Using 2001 data, the report estimates the percentage decrease in IHD and stroke burden that could be expected if population exposure to a risk factor were reduced to zero by calculating the

population attributable fraction for each of the key CVD risk factors. The report found that hypertension, high cholesterol, overweight and obesity, smoking, low fruit and vegetable intake, and physical inactivity were the leading contributors to IHD and stroke burden worldwide (54).

2.6 Major proximal risk factors for CVD

This section described the major risk factors for CVD in more detail. The section begins with behavioral risk factors, including tobacco use, dietary factors, alcohol, and physical activity. This is followed by the major biological risk factors that mediate the role of these behaviors in leading to CVD including obesity, blood pressure, blood lipids, and diabetes. Finally, additional contributing factors are also discussed, including genetics and gender difference.

Tobacco

In the Global Burden of Disease study, Lopez et al. (2006) estimated that in 2000, 880,000 deaths from CHD and 412,000 deaths from stroke were attributable to tobacco (10). These data are based on updated estimates of the relative risk of death among smokers for CHD, stroke, and hypertensive heart disease. The relative risks are highest in young people. However, the most common type of tobacco-related CVD deaths varies around the world. In India, a higher proportion of smokers die from CHD; in China, tobacco kills more through stroke (55).

Smoking cessation has been shown to have significant impacts on reducing CHD. Smoking cessation leads to significantly lower rates of reinfarction within 1 year among patients who have had a heart attack and reduces the risk of sudden cardiac death among patients with CHD. There is consensus in the literature that CVD risk drops precipitously within the first 2 to 3 years of smoking cessation. Although the specific timeline of risk reduction depends on the number of years as a smoker and the quantity of tobacco smoked daily, it is conceivable that, over time, former smoker's CVD risk can drop to levels similar to that of someone who has never smoked (56). Two major trends are of real concern with respect to the future of tobacco-related CVD. First, in most parts of the

world, the smoking rates are higher among the poorest populations. The second worrisome trend is in smoking among girls. In most parts of the developing world, women smoke at a significantly lower rate than men, a disparity that could help explain the lower rates of cardiovascular mortality among women (57). If future generations of girls catch up to boys and smoke at the rates that men do today, CVD and associated tobacco-related death rates will rise sharply.

Dietary Factors

The relationship between CVD and diet is one of the most studied relationships in epidemiology. Several key relationships identified decades ago remain valid, while others have evolved in the light of better-quality research. WHO and FAO reviewed the evidence on the relationship between diet, physical activity, and CVD in the context of a broader review of the impact on all chronic diseases. High dietary intakes of saturated fat, trans-fats and salt; low intake of fruits, vegetables and fish are linked to cardiovascular risk. Approximately 16 million (1.0 per cent) DALYs and 1.7 million (2.8 per cent) of deaths worldwide are attributable to low fruit and vegetable consumption (58). Elimination of trans-fat and replacement of saturated with polyunsaturated vegetable oils lowers coronary heart disease risk (58). A healthy diet can contribute to a healthy body weight, a desirable lipid profile and a desirable blood pressure, leading to reduced risk of CVD.

Oils

The rapid rise in the production and consumption of tropical oils has worried many CVD researchers because of their adverse effects on CVD risk. Healthy oils are those that contain no commercially introduced trans fatty acids, are low in saturated fatty acids, and are high in mono- and polyunsaturated fatty acids. Nutritionally, the most important mono- and polyunsaturated fatty acids are oleic acid and linoleic acid, respectively. Olive and canola oils have high concentrations of oleic acid, whereas nonhydrogenated soybean oils and sunflower oils have high concentrations of linoleic acid. All four of these oils are also low in saturated and trans fats, but their shelf lives and cooking properties (smoke point, flavor, etc.) vary. High and mid-oleic sunflower oils both have long shelf lives, but

unfortunately they remain relatively expensive and less abundant in many low and middle income countries. In order to be truly effective, low and middle income countries that have high levels of oil consumption will need to develop affordable supplies of healthy oils at prices that are competitive with tropical oils. Transitioning from less healthy tropical oils to more healthy oils could significantly reduce the amount of saturated and trans-fatty acids used in highly processed foods and daily cooking.

Salt

There is a strong and robust base of evidence that excessive sodium intake significantly increases CVD risk and that reduction in sodium intake on a population level decreases CVD burden (59). The most well-established mechanism by which sodium intake increases CVD risk is by increasing blood pressure. Numerous studies have found that there is a continuous and graded relationship between salt intake and blood pressure. This relationship has been confirmed in epidemiological, animal, population, migration, intervention, and genetic studies. Furthermore, population studies have established that reductions in sodium intake lead to declines in systolic and diastolic blood pressure, which in turn leads to a decrease in heart attacks and strokes. For example, since the 1970s, salt intake in Finland has been reduced by approximately one-third. This has led to a reduction in systolic and diastolic blood pressure (BP) by more than 10 mmHg, a pronounced decrease of 75–80% in both stroke and CHD mortality and a remarkable increase of 5–6 years in life expectancy (60). In their recent major review of sodium trends and impact, H E and MacGregor concluded that a reduction in salt from the current global intake of 9 to 12 g/day to the recommended levels of 5 g/day would have a major impact on BP and on CVD (58). Salt's impact on CVD, however, extends beyond blood pressure. Animal and epidemiological studies have found that a diet high in sodium may directly increase risk of stroke, which is independent and additive to salt's effect on BP (59).

Alcohol

The global burden of diseases attributable to alcohol has recently been summarized, leading to the conclusion that alcohol is one the largest avoidable risk factors in low and

middle income countries. It has long been known that excessive alcohol intake is associated with increased risk for hypertension, stroke, coronary artery disease, and other forms of CVD; however, there is also a robust body of evidence in a range of populations that suggests that light to moderate intake of alcohol may reduce the risk of CHD. Indeed, research suggests that the relationship between alcohol intake and CVD outcomes follows a “J” curve, with the lowest rates being associated with low to moderate intakes of alcohol (61). It is important to recognize that, as with any discussion of alcohol and health, the key issues are the quantity of alcohol consumed and the risk or benefit conferred by consumption. Although evidence indicates that low to moderate alcohol use can reduce the risk of CHD, excessive and harmful use clearly increases CVD risk. Alcohol may also contribute to overweight and obesity as it is a significant source of daily calories in many countries (62). It is also important to consider the demonstrated negative health effects of excessive and harmful alcohol use on other diseases such as neuropsychiatric disorders, cirrhosis of the liver, and various cancers.

Physical Activity

WHO highlighted the importance of physical activity as a key determinant of obesity, CVD and diabetes. For decades, evidence of the relationship between physical activity and CVD, independent of effects on weight and obesity, has strengthened. Increasing physical activity—including brisk walking—has been shown to decrease the risk of chronic diseases such as CHD, stroke, some cancers (e.g., colorectal and breast cancer), type 2 diabetes, osteoporosis, high blood pressure, and high cholesterol. Physical activity is also important for weight control and maintenance. In addition, regular physical activity is associated with a decreased risk of depression and improved cognitive function. Moreover, people who are physically active have improved quality of life and reduced risk of premature death (63).

Guthold et al. (2008) recently published new data on levels of physical inactivity in 51 countries, most of which were low or middle income, and observed several trends. Globally, with the exception of several Eastern European countries (Croatia, the Czech Republic, Hungary, Kazakhstan, the Russian Federation, Slovenia, and the Ukraine),

women were more likely to be physically inactive than men. Further, adults over 50 years of age were more likely to be inactive than younger adults, and city dwellers were more likely to be inactive than those who lived in rural areas. Physical inactivity levels were, with a few exceptions, similar in Eastern European, South Asian, and Western Pacific countries. In most of these countries, between 5 and 10 percent of men and between 10 and 16 percent of women were found to be physically inactive. By contrast, there was considerable variation in the levels of physical activity in both men and women within and across African, American, and Eastern European countries. For example, while women in 7 of the 18 African countries surveyed had the lowest levels of physical inactivity (fewer than 10 percent classified as physically inactive), Guthold found that more than 40 percent of women in Namibia, Swaziland, and South Africa were physically inactive. Despite the heterogeneity of the data, the study indicated that levels of physical inactivity in a number of low and middle income countries and among certain subgroups, particularly women aged 60–69 years, are disconcertingly high (64).

Overweight and Obesity

WHO and FAO reviewed the evidence on the relationship between obesity and the risk of CVD and concluded that overweight and obesity confer a significantly elevated risk of CHD (65). Increased body mass index (BMI) is also associated with greater risk of stroke in both Asian and Western populations (65). The association between obesity and CVD is partly, but not completely, mediated through hypertension, high cholesterol, and diabetes. Abdominal or central obesity measured by waist-to-hip ratio or waist circumference is associated with both CHD and stroke independent of BMI and other cardiovascular risk factors. Moreover, obesity is also an independent risk factor for other cardiovascular outcomes, such as congestive heart failure and sudden cardiac death.

Excess energy intake is one of the key contributors to obesity. As highlighted earlier, the lack of data limits policy makers' abilities to focus attention on which dietary components lend themselves to effective interventions that would reduce total calorie intake. In those countries that do have data, the collection methods vary so direct comparisons are not possible; however, a review of the data does indicate that the dietary contributors to total energy intake vary by country. National surveys of calorie intake

from India indicate that in urban areas, cereals account for 56 percent of intake, compared to about 9 percent each for edible oils and dairy, 1 percent for meat and fish, and 0.4 percent for all beverages (66). In China, cereals also dominate and account for 58 percent of total calorie intake compared to meat (13 percent) and cooking oils (17 percent) (67). In some developing countries, consumption of sugar-sweetened beverages has increased dramatically in recent decades. In Mexico, for example, it is estimated that adolescents consume more than 20 percent of their total energy intake from caloric beverages (68). Because of its excess caloric and sugar content, increasing consumption of sugar-sweetened beverages may have important implications for obesity and cardiometabolic risk. Maintaining the relatively low per capita consumption of sugar-sweetened beverages in countries like India and China is a potential target of prevention programs. In India, all beverages account for less than 0.5 percent of total calories (66). The equivalent figure in the United Kingdom is about 16 percent for all beverages for young adult men between 19 and 24 years of age with sweetened soft drinks accounting for about a third and alcohol the remainder (69).

Hypertension

A recent review of the global burden of high blood pressure found that approximately 54 percent of stroke, 47 percent of IHD, 75 percent of hypertensive disease, and 25 percent of other CVDs were attributable to hypertension. This equates to an annual burden of approximately 7.6 million deaths, or 13.5 percent of the total number of annual global deaths, attributable to high blood pressure (70). Furthermore, Lawes et al. (2008) found that more than 80 percent of the attributable burden of hypertension in 2001 occurred in low and middle income countries, and both another recent review and an analysis commissioned for this report found the prevalence of hypertension to be equally high in developed and developing countries (71).

In Sub-Saharan Africa, hypertension is a predominant driver of CVD. Hypertensive heart disease and stroke, rather than ischemic heart disease, account for the majority of the CVD burden in the region, especially among black Africans (72).

Blood Lipids

Researchers have studied the role of blood lipids in the development of atherosclerosis and the increase of CVD risk for decades. The Framingham Study first demonstrated the link between hypercholesterolemia and increased risk of CHD in the 1960s with the finding that lower levels of high-density lipoprotein (HDL) cholesterol as well as elevated levels of low-density lipoprotein (LDL) cholesterol were associated with increased CHD risk (73). Subsequent studies confirmed these results and further established that elevated triglycerides also increase CVD risk (74). Furthermore, randomized controlled trials have shown that reduction of LDL cholesterol, both in primary and secondary prevention, is associated with reduced coronary event rates (75). Reductions in LDL cholesterol have also been associated with a lowered incidence of stroke, although the data are not as strong as for CHD (76). The INTERHEART study recently confirmed that there was a graded relationship between abnormal lipid levels and risk for CHD in all regions of the world. In fact, the INTERHEART study found that abnormal blood lipids were the most important risk factor for myocardial infarction by odds ratio in all global regions (77).

Diabetes

Around the world, diabetes is growing increasingly common and is a significant contributor to CVD risk. People with diabetes have a more than two-fold greater risk of fatal and nonfatal CVD compared to non-diabetics, with some indication that diabetes mellitus may confer an equivalent risk of having had a cardiovascular event (78). In fact, CVD is the leading cause of morbidity and mortality in people with diabetes (79).

The magnitude of the risk of CVD associated with diabetes is even greater in women and younger individuals. Indeed, there is substantial evidence that diabetes mellitus may erase, or substantially attenuate, the “female advantage” in the risk of CVD observed in non-diabetics, and that having diabetes may be equivalent to aging by at least 15 years with regard to the clinical manifestations of CVD (80).

Diabetes is emerging as a particular concern in Asia, where more than 110 million individuals were living with diabetes in 2007, a large proportion of whom were young and middle aged. Asians tend to develop diabetes at a relatively young age and low BMI,

and by 2025 the number of individuals with diabetes in the region is expected to rise to almost 180 million, of which approximately 70 million will be in India and almost 60 million in China. The reasons for this increased risk are still being fully elucidated; however, “normal weight” Asians often exhibit features of abdominal or central obesity, which is particularly detrimental to insulin resistance and glucose metabolism. Moreover, the increased risk of gestational diabetes combined with exposure to poor nutrition in utero and over-nutrition in later life may contribute to increased diabetes, resulting in a situation of “diabetes begetting diabetes” (81).

The balance of risks and benefits associated with intensive glucose control has been assessed in recent clinical trials, which have convincingly demonstrated beneficial microvascular outcomes of diabetes. By contrast, these trials have individually failed to show such an effect on cardiovascular outcomes. However, the extension of the follow-up of the Diabetes Control and Complications Trial in type 1 diabetes and the United Kingdom Prospective Diabetes Study in type 2 diabetes have shown that intensive glucose control substantially lowered the risk of cardiovascular outcomes (82,83). Recently conducted meta-analyses of relevant trials in people with type 2 diabetes have also consistently shown that intensive glucose control reduces the risk of major cardiovascular events by approximately 10 percent, primarily driven by a 10 to 15 percent reduction in the risk of CHD, compared with standard treatment in people with diabetes. Interestingly, this benefit appeared to be independent of concurring cardiovascular risk factors (84).

Genetics

Researchers have recognized for decades that family history of CVD is associated with increased atherosclerotic risk of heart disease, which led to the presumption of a genetic component to CVD. There are several well-characterized single-gene disorders that contribute to CVD, such as certain forms of familial hypercholesterolemia linked to mutations of the apolipoprotein B gene, and during the past few years, there have been major advances in the identification of genetic risk factors for CHD, stroke, and CVD risk factors such as blood pressure, blood lipids, obesity, and diabetes (85). The identification of genetic loci associated with CVD, such as 9p21, has led to major

advances in understanding the pathophysiology of CVD (86). The prevailing view within the research community is that the genetic underpinnings of most common forms of CVD involve a complex interplay of many different genes, and much work remains to develop a more thorough understanding of the complex gene–gene and gene–environment interactions involved in the development of CVD (87).

Indeed, in addition to the investigation of genes that influence CVD and its risk factors, there has recently been a surge in research examining how environmental factors affect gene expression. Although research indicates that gene expression is most sensitive to environmental influence from conception to early life, there is also evidence that environmentally related gene expression changes can occur throughout life (88). This is an important emerging area of research for CVD. Future findings could have implications to help elucidate the physiological processes by which individuals with similar CVD risk profiles have different outcomes. Future research also could conceivably help develop new prevention and treatment strategies aimed at taking advantage of exogenous mechanisms that enhance or suppress the expression of key genes that play a role in mediating the development of CVD.

Gender Differences in CVD Risk

Although CVD has sometimes been considered a disease that predominantly affects men, it is the leading cause of death among both men and women globally (89). There are, however, a number of notable gender differences in CVD incidence, mortality, risk-factor profiles, outcomes, and clinical presentation. These differences remain consistent across populations and regions and are thus important to consider when developing CVD prevention and treatment programs.

In all but the oldest age groups, CVD prevalence, incidence, and mortality rates tend to be higher for men than for women. This finding has remained consistent historically and across countries and regions (90). In addition, women experience their first cardiovascular events later in life than men. The INTERHEART study found that, on average, women experience their first MI 9 years later than men (91).

The reason most often cited for these gender differences is a protective effect of estrogen on the development of CVD risk factors, most notably hypertension and dyslipidemia (92). Estrogen is thought to contribute to premenopausal women's tendency to have lower systolic blood pressure, higher levels of HDL cholesterol, and lower triglyceride levels than men (93). The specific mechanisms of this protection have not been fully elucidated; however, estrogen is known to affect the atherosclerotic and blood-lipid control process in a number of different ways (94). The erosion of this protection that occurs after menopause provides further evidence of estrogen's protective role. Indeed, by age 75, women tend to have higher rates of hypertension and CVD than men (95). Risk of stroke is similar for men and women (35).

CHAPTER - 3
SUBJECTS AND METHODS

Chapter 3

Subjects and Methods

WHO STEPS chronic disease risk factor surveillance guideline was followed in this study. As resource was limited, only first two steps i.e. questionnaire and physical measurements was done for this study.

3.1: Study design: It was a cross sectional survey focused on the prevalence of risk factors of cardiovascular disease among adults in a slum area of Dhaka city.

3.2: Study period: Study duration was one year (from July 2014 to June 2015). Protocol writing took first 2 months. Preparatory activities including questionnaire development, training and piloting were done for 2 month. Data collection period was 4 months. Rest of the time was spent for data processing, analysis and report writing. Besides literature were reviewed during the whole study period.

3.3: Study area: The study was conducted at Agargaon slum area and this area was selected purposively.

3.4: Study population: The study population for this study included slum men and women aged 35 years and above living in Agargaon, Sher-E-Bangla Nagar, Dhaka. In general, the target population of the study may be included individuals residing in all geographic slum areas of the Dhaka city.

3.5: Study sample: Individual adult (may be more than one) from all household (A dwelling in which persons either related or unrelated living together and taking food from the same kitchen) as there was no sampling frame.

3.6: Inclusion criteria

All slum men and women aged 35 years and above living in the slum

3.7: Exclusion criteria

- Not willing to participate in the study
- Having mental disorders
- Pregnant women

3.8: Sample size

A representative sample size of 461 was taken for the study. This sample size has been calculated using the following formula:

$$n = \frac{Z^2 pq}{e^2}$$

Where, $Z=1.96$

p = prevalence of diabetes was 10% (lowest prevalence among all CVD risk factors (31).

$q (1-p) = 0.9$

e (precision level/ margin of error) = 0.05

Initial sample calculation = $3.8416 * 0.9 * 0.1 / 0.05 * 0.05 = 138$

After multiplying by the design effect and number of age-sex estimates:

$n = 138 * 1.5 * 2 = 415$

As the resources were extremely limited, age-sex estimates were obtained only for the entire age span of the survey. This was 2 for male and female sexes.

Now, the **final sample size** was obtained by adjusting for expected non-response:

$n = 415 / 0.9 = 461$.

3.9: Sampling technique: Non probability convenient sampling was used for data collection on the basis of inclusion and exclusion criteria.

3.10: Data collection techniques/methods

Face to face interview was done by using structured questionnaire. Informed consent was taken prior interview and whole procedure was described prior to start filling of questionnaire. The interviews and measurements were done at household level.

Training for data collectors for 3 days long



Meeting with slum leaders



Explaining the importance of this study



Informed consent obtained



Data collection started door to door

Along with socio-demographic information, behavioral habit, information on hypertension and diabetes mellitus, anthropometric measurement (height, weight, waist circumference) as well as measurement of blood pressure was done.

Anthropometric measurements:

Height, weight and waist circumference was measured to calculate their body mass index (BMI), thereby obesity. The standing height was measured with a stadiometer with minimal cloths. A stadiometer consist of a metric tape affixed to a vertical surface and a movable headpiece attached to the vertical surface that can be brought down to the crown of the head. Participants will be asked to remove their footwear and head gear, if any, and stand on the board facing the interviewer, feet together, heels against the board, knees straight. Looking straight ahead and not to tilt head up and making sure that eyes are at the same level as the ear. The subjects were asked to inhale deeply and maintain a fully erect position. The headpieces were brought down until it touches the head; sufficient pressure was applied to compress the hair. Three measurements were taken three times and if the difference among reading found to be less than 1 cm, the mean measurement

was taken and recorded to the nearest 0.1 cm. If the reading falls between two values, the lower reading was recorded.

The body weight was measured using a platform beam scale. The beam of the platform scale was graduated so that it can be read from both sides; the calibration of the scale was done before taking the weight. The subjects stand still over the center of the platform with body weight evenly distributed between both feet with light indoor clothing. Weight was recorded to the nearest 0.1 kg.

Waist circumference was measured by plastic measuring tape, maintaining privacy of the participants; inside room or area screened off from other people. This was measured directly on the skin at the end of a normal expiration with the arms relaxed at the sides, at the midpoint between the lower margin of the last palpable rib and the top of the hip bone. Waist circumference was measured to nearest millimeter.

Blood pressure measurement

Blood pressure was measured in sitting position; with calf at the level of the heart. After 5 minute of rest a second reading was taken. If the difference between two readings was more than 5 mm of Hg, a third reading was taken.

3.11: Data collection Tools

- Structured questionnaire- The questionnaire was developed with minor adaptation of WHO STEP wise Surveillance (STEPS) questionnaire. All the core variables along with some expanded variables from step 1 were incorporated. From step 2 physical measurements (height, weight, waist circumference and blood pressure) was included.
- BP machine and stethoscope
- Physical measurements tools (weight machine, height measuring scale, waist measuring non stretch tape)
- Pictorial show cards or measuring cups of different sizes.

3.12: Data analysis

Version 17 of SPSS was used for data entry and analysis. Obtained information was presented in the form of tables and graphs. Descriptive statistics such as mean, SD, frequency, percentage was used. Chi-square test was done to see association. Binomial logistic regression was conducted to identify the effect of risk factors on dependent variables.

3.13: List of variables

3.13.1: Demographic variables

1. Age
2. Sex
3. Education
4. Occupation
5. Religion
6. Marital status
7. Family size
8. Monthly family income
9. House type

3.13.2: Behavioral variables

1. Tobacco use
2. Unhealthy diet
 - 2.1 Low fruit and vegetable consumption
 - 2.2 High salt and oil consumption
3. Physical activity
4. Harmful alcohol consumption


3.13.3: Biologic variables

1. Overweight and obesity
2. Waist circumference
3. High blood pressure
4. Diabetes

3.13.4: Non modifiable variable

1. Family history of heart attack or stroke*

3.14: Conceptual Framework



3.15: Operational Definitions

3.15.1: Tobacco use:

Information of tobacco use was collected for both smoking and smokeless forms.

Cut-offs for tobacco use

The two main tobacco indicators that are associated with an increased risk of developing chronic diseases including cardiovascular diseases are:

- Current smoker and
- Daily smoker

The smoking status of the participants was categorized into 'current smoker' those who have smoked tobacco in past 30 days and 'daily smoker' those who smoke any tobacco products every day (6). They were asked about the initiation and duration of use by means of age and year(s) both. Frequency of use of products was also asked. Question was asked to non smokers if he or she were exposed to second hand smoke in last seven days at their home or workplace.

Smoking forms of tobacco includes biri, cigarette (both manufactured and roll your own-RYO), pipe, cigars, water pipe also known as shisha, hookah or hubble-bubble (6).

Smokeless tobacco includes jarda, sada pata, snuff like gul, betel nut, chewing tobacco e.g., plug, loose-leaf, chimo, toombak, guthha or twist, betel leaf etc (6). Show cards were used for different forms of tobacco.

3.15.2: Low fruit and vegetable consumption

Minimum daily total requirement of fruit and vegetable was 5 servings of each. One standard serving size equals to 80 grams (6). Servings were measured by showing the pictorial show cards or measuring cups.

- For raw green leafy vegetables, 1 serving = one cup;
- For cooked or chopped vegetables, 1 serving = ½ cup;
- For fruit (apple, banana, orange), 1 serving = 1 medium size piece;
- For chopped, cooked and canned fruit, 1 serving = ½ cup; and
- For juice from fruit, 1 serving = ½ cup

3.15.3: High salt and oil consumption

Salt: The recommended level of salt intake is 5 grams/day (21).

Oil: For our country, the recommended amount of edible oil to be consumed per day is 5 tsp or 25 ml for men and 4 tsp or 20 ml for women (96).

3.15.4: Physical activity

Physical activity was measured according to Global Physical Activity Questionnaire method (97). The Global Physical Activity Questionnaire (GPAQ) was developed by WHO for physical activity surveillance in countries. It collects information on physical activity in three domains. These domains are:

- Activity at work
- Travel to and from places
- Recreational activities

Physical activity was defined as any bodily movement produced by skeletal muscles that require energy expenditure. Physical activity was categorized into vigorous, moderate and sedentary activity. A vigorous-intensity activity was defined as any activity that causes large increase in breathing or heart rate, if continued for at least 10 minutes (e.g. running, carrying or lifting heavy loads, digging or construction work). Moderate-intensity activity was defined as any activity that causes small increase in breathing or heart rate, if continued for at least 10 minutes (brisk walking or carrying light loads).

Physical activity related to work means whether the work involves vigorous-intensity activity (running, carrying or lifting heavy loads, digging or construction work) or moderate-intensity activity (brisk walking or carrying light loads). Physical activity related to travel to and from places (transportation) means walking or using a bicycle (*pedal cycle*) for at least 10 minutes continuously to get to and from places. Physical activity related to recreational activities means sports, fitness or recreational activities like running or playing football, volleyball, brisk walking, cycling, swimming and can be of vigorous-intensity or moderate-intensity. Continuous activity of at least ten minutes was taken into account.

Physical activities was measured in the survey by asking the respondents about their weekly and daily vigorous and moderate activities during work and leisure time, activities related with transport and time spent in sedentary position. Respondents were asked for number of days they do vigorous and moderate activities in a typical week. In such a day, how much time they spent (in minutes) in these type of activities. Similarly they were asked about time spent on transport related activities. All type of physical activities was transferred in minutes per day. Then the total duration was converted into metabolic equivalents (MET minutes/week). METs were commonly used to express the intensity of physical activities. MET is the ratio of a person's working metabolic rate relative to their resting metabolic rate. One MET was defined as the energy cost of sitting quietly and is equivalent to a caloric consumption of 1 kcal/kg/hour. It is estimated that compared with sitting quietly, a person's caloric consumption is three to six times higher when being moderately active (3-6 METs) and more than six times higher when being vigorously active (>6 METs). However for our study we took four times for moderate and eight times for vigorous-intensity activity than sedentary position (sitting quietly). MET-minute was calculated according to the STEPS protocol as follows: one minute in sedentary position (sitting quietly) equal to 1 MET-minute. One minute in moderate and transport related activities equal to 4 MET-minutes and one minute in vigorous activities equal to 8 MET-minutes. MET values were applied only to vigorous and moderate intensity variables in the work and recreation settings and to transport related activities. All transport related activities were considered as moderate intensity activities. All MET-

minutes for different forms of physical activities was added together to get total physical activities in MET-minutes. Then physical activities were categorized in high, moderate and low type. Those who spent 3000 or more MET-minutes per week, they were categorized as high physical activity group. Those who spend between 600 up to 3000 MET-minutes, they were categorized as moderate physical activity. For those who spent less than 600 MET-minutes, they fall in low activity category.

3.15.5: Alcohol consumption

Alcohol consumption was measured by asking the respondents if they consumed ever, within past 12 months, and within past 30 days. They were also asked about the maximum frequency of "standard" alcoholic drinking in a single occasion in past 12 months and in past 30 days. They were also asked about daily and weekly standard alcoholic drink. One standard drink equals to 10 grams of ethanol and was measured by showing the pictorial show cards of different size drinking glasses (shown in annexure). The terms to be used were listed as follows:

- **Current drinker:** who drank within past 30 days.
- **Daily drinker:** who drink daily.
- **Alcohol consumer:** who drank in past 12 months.
- **Binge drinkers:** 5 or more drinks on a single occasion for men or 4 or more drinks on a single occasion for women, generally within about 2 hours (98).
- **Heavy drinking:** For men, heavy drinking was typically defined as consuming 15 drinks or more per week. For women, heavy drinking was typically defined as consuming 8 drinks or more per week (99).
- **Moderate drinking:** According to the Dietary Guidelines for Americans, moderate alcohol consumption was defined as having up to 1 drink per day for women and up to 2 drinks per day for men (100).

3.15.6: Body mass index: Body mass index was calculated from the body weight and height of the subjects using the following formula: BMI= weight in kg/ height in m². BMI was interpreted as follows:

<u>BMI</u>	<u>Nutritional class</u>
<16	Severe malnutrition
16-16.9	Moderate malnutrition
17-18.4	Mild malnutrition
18.5-24.9	Normal
25-26.9	Overweight
27-29.9	Obesity
30-34.9	Moderate obesity
35-39.9	Severe obesity
40 or more	Morbid obesity

3.15.7: Waist circumference: Increased when it was ≥ 94 cm in men and ≥ 80 cm in women (6).

3.15.8: Hypertension

A person was considered being mildly hypertensive if the systolic value (SBP) ≥ 140 mmHg and/or the diastolic value (DBP) ≥ 90 mmHg. Moderate hypertension has been defined as SBP ≥ 160 mmHg and/or DBP ≥ 100 mmHg (101).

Normal	SBP 90-119 and 60-79
Pre-hypertension	SBP 120-139 or DBP 80-89
Stage 1 HTN	SBP 140-159 or DBP 90-99
Stage 2 HTN	SBP ≥ 160 or DBP ≥ 100

3.15.9: Diabetes

A person was considered having diabetes if she/he was previously diagnosed by a physician as diabetic and was under treatment or not.

3.15.10: Socio-economic status

Socioeconomic status of study subjects could not be classified. So instead of low socioeconomic status, I consider low family monthly income as risk factor for CVD. Family monthly income 10000 taka or less was considered as risk factor.

3.16: Ethical Consideration

Permission from ethical review committee of Bangladesh Institute of Health Science was taken. Informed consent was taken before starting the interview. Risk status of the respondents was provided which help them to take appropriate preventive measures. There was no human risk as there was no invasive procedure. The study did not divulge the name and personal information of any individual respondent at any point of time. Respondent's right to refuse and withdraw from the study was accepted.

CHAPTER 4
RESULTS

Chapter 4

Results

This cross-sectional study was conducted with the aim to estimate the prevalence of cardiovascular disease risk factors among slum population of Dhaka city. Though hypertension is a major modifiable risk factor for CVD, it is also included as one of the cardiovascular disease. In this study, hypertension was considered as the primary outcome and association of hypertension with various socio-demographic characteristics, modifiable and non-modifiable risk factors were sought. The result of the study has been described in five major sections as follows:

- 1. Socio-demographic characteristics and risk factors distribution among male and females**
- 2. Association between hypertension and various socio-demographic characteristics, modifiable and non-modifiable risk factors**
- 3. Effect of various modifiable and non-modifiable risk factors on hypertension**
- 4. Effect of non-modifiable risk factors on various modifiable risk factors**

Table 4.1.1: Socio-demographic characteristics of study subjects- age, marital status, education, occupation, economic status, religion, family size & house type

Variables	Total n=461(100)	Male (n=241)	Female (n=220)	p value
Age in years				
35 to < 45	285 (61.8)	135 (56.0)	150 (68.2)	0.001
45 to <55	111 (24.1)	57 (23.7)	54 (24.5)	
55 to <65	47 (10.2)	34 (14.1)	13(5.9)	
≥65	18(3.9)	15(6.2)	3(1.4)	
Mean (±SD) age	43.04 ±8.86	45.03±9.47	40.85±7.57	0.000
Marital status				
Married	433 (93.9)	230(95.4)	203(92.3)	0.009
Unmarried	13(2.8)	9(3.7)	4(1.8)	
Divorced	7(1.5)	0(0.0)	7(3.2)	
Widower/widow	8(1.7)	2(0.8)	6(2.7)	
Education				
No formal education	263(57.0)	127(52.7)	136(61.8)	0.013
Primary not completed	83(18.0)	37(15.4)	46(20.9)	
Primary	87(18.9)	57(23.7)	30(13.6)	
Secondary	22(4.8)	16(6.6)	6(2.7)	
Higher secondary	5(1.1)	3(1.2)	2(0.9)	
Occupation				
House hold work	181(39.3)	1(0.4)	180(81.8)	0.000
Small business	97(21.0)	79(32.8)	18(8.2)	
Day labourer	55(11.9)	51(21.2)	4(1.8)	
Driver	29(6.3)	29(12.0)	0(0.0)	
Rickshaw puller	18(3.9)	18(7.5)	0(0.0)	
Private employee	17(3.7)	11(4.6)	6(2.7)	
Industrial labourer	15(3.3)	10(4.1)	5(2.3)	
Family income/month				
≤10000	169(36.7)	85(35.3)	84(38.2)	0.234
10001 - 15000	183(39.7)	90(37.3)	93(42.3)	
15001– 20000	89(19.3)	55(22.8)	34(15.5)	
>20000	20(4.3)	11(4.6)	9(4.1)	
Mean (±SD)	13754.88±4592.59	14026.97±4771.25	13456.81±4379.97	0.183
Religion				
Islam	445(96.5)	230(95.4)	215(97.7)	0.389
Hindu	12(2.6)	8(3.3)	4(1.8)	
Christian	4(0.9)	3(1.2)	1(0.5)	
Family member				
≤2	10(2.2)	8(3.3)	2(0.9)	0.199
3-4	177(38.4)	90(37.3)	87(39.5)	
≥5	274(59.4)	143(59.3)	131(59.5)	
Type of house				
Tin shade	460(99.8)	240(99.6)	220(100)	0.339
Brick build	1(0.2)	1(0.4)	0(0.0)	

*Values expressed as n(%) in parentheses or M±SD, as appropriate; M, mean; SD, standard deviation, p value, significance between male and female participants; p value was obtained by χ^2 test for categorical variables & t- test for continuous variable; 0.05 was considered as level of significance.

Total 461 adult slum people were studied, of them males were 241(52.3%) and females were 220(47.7%). Male female proportion was almost same. Table 4.1.1 shows that mean age of the respondents was 43.04 years with standard deviation of 8.86 years whereas the mean age of male and female were 45.03 (SD±9.47) and 40.85 (SD±7.57) years respectively and the male female distribution was statistically different ($p < 0.05$). The age range of the study subjects was from 35 to above. Out of 461 respondents, majority of the study subjects (62%) was in the age range of 35 up to 45 years. Senior citizens (55 years or more) were only about 14% of study subjects. The lower age range study subjects were significantly higher than higher age range subjects ($p < 0.05$). Almost all of the respondents (93.9%) were married where male and female was 95.4% and 92.3% respectively. More than half of study subjects (57%) had no formal education where male and female distribution was 52.7% and 61.8% respectively. Only 24.8% subjects completed more than primary level of education and the literacy rate was significantly higher in male than ($p < 0.05$). Majority of the females (81.8%) were house hold worker, compared to a very negligible (0.4%) male. About one third (32.8%) males were small businessmen, compared to 8.2% females. Day laborer was about 12%, where male and female was 21.2% and 1.8% respectively. Driver and rickshaw pullers were all males, 12% and 7.5% respectively. The mean monthly family income of the respondents was 13754.88 (SD±4592) whereas for male and female it was 14026.97 (SD±4771.25) and 13456.81 (SD±4379.97). A little more than three-fourth (76.4%) of the respondent's monthly family income were up to 15000 BDT. A little less than one fourth (23.6%) of the respondent's monthly family income was above 15000 taka. Monthly family income did not vary statistically among male and female ($p > 0.05$). Almost all of the respondents (96.5%) were muslim, the rates of male and female were 95.4% and 97.7% respectively. About 60% of the respondents had a family size of 5 or more persons. Almost 100% of the respondents lived in tin shed house.

Table 4.1.2: Behavioral characteristics of the respondents

Variables	Total n=461(100)	Male (n=241)	Female (n=220)	<i>p value</i>
Smoking tobacco				
Current smoker	181(39.3)	177(73.4)	4(1.8)	0.000
Non smoker	280(60.7)	64(26.6)	216(98.2)	
Daily smoker	175(38.0)	173(71.8)	2(0.9)	0.000
Mean (\pm SD) age of starting smoking	16.95 \pm 4.47	16.83 \pm 4.5	23.66 \pm 10.96	0.394
Mean (\pm SD) daily consumption of sticks	14.91 \pm 5.82	15 \pm 5.8	11.25 \pm 6.29	0.204
Types of tobacco products (n=181, male-177, female-4)				
Cigarette	164(90.6)	162(91.5)	2(50.0)	0.014
Biri	16(8.8)	14(7.9)	2(50.0)	
Hand rolled cigarette	1(0.6)	1(0.6)	0(0.0)	
Past daily smoker among non-daily smoker (n=286, male-68, female-218)	30(10.5)	29(42.6)	1(0.5)	0.000
Non smokers who exposed to second hand smoke at home (n=280, male-64, female-216)	175(62.5)	26(40.6)	149(69.0)	0.000
Non smokers who exposed to second hand smoke at workplace (n=280, male-64, female-216)	35(12.5)	16(25.0)	19(8.8)	0.001
Smokeless tobacco				
Current smokeless tobacco user	219(47.5)	87(36.1)	132(60.0)	0.000
Daily smokeless tobacco user (n=219, male-87, female-132)	188(85.8)	69(79.3)	119(90.2)	0.024
Types of smokeless tobacco (n=219, male-87, female-132)				
Zorda	175(79.9)	73(83.9)	102(77.3)	0.284
Sada pata	23(10.5)	9(10.3)	14(10.6)	
Guul	21(9.6)	5(5.7)	16(12.1)	
Mean (\pm SD) frequency of daily smokeless tobacco use	8.36 \pm 4.87	8.20 \pm 5.56	8.45 \pm 4.43	0.204
Physical activity				
Low physical activity	167(36.2)	79(32.8)	88(40.0)	
Moderate physical activity	151(32.8)	57(23.7)	94(42.7)	0.000
High physical activity	143(31.0)	105(43.6)	38(17.3)	
Work related vigorous activity	98(21.3)	90(37.3)	8(3.6)	0.000
Travel to and from places by bicycle/walking	213(46.2)	123(51.0)	90(40.9)	0.059
Vigorous work during recreational activities	4(0.9)	3(1.2)	1(0.5)	0.361
Mean(\pm SD) minutes of sedentary activity in a day	159.76 \pm 71.28	163.44 \pm 76.67	155.73 \pm 64.80	0.243

Variables	Total n=461(100)	Male (n=241)	Female (n=220)	<i>p value</i>
Alcohol consumption				
Lifetime abstainer of alcohol	413(89.6)	196(81.3)	217(98.6)	0.000
Alcohol consumer	16(3.5)	16(6.6)	0(0.0)	
Current drinker	8(1.7)	8(3.3)	0(0.0)	
Daily drinker	4(0.9)	4(1.7)	0(0.0)	
Binge drinker	4(0.9)	4(1.7)	0(0.0)	
Heavy drinker	2(0.4)	2(0.8)	0(0.0)	
Moderate drinker	2(0.4)	2(0.8)	0(0.0)	

**Values expressed as numbers and percentages in parentheses or M±SD, as appropriate; M, mean; SD, standard deviation, p value, significance between male and female participants; p value was obtained by χ^2 test for categorical variables & t- test for continuous variable; 0.05 was considered as level of significance*

Table 4.1.2 shows that overall prevalence of smoking was 39.3%, it was 73.4% among men and 1.8% among women, prevalence found significantly higher among male ($p < 0.05$). Overall prevalence of daily smoking was 38%, the rates were 71.8% and 0.9% among male and female respectively. Daily smoking rate was found significantly higher among male ($p < 0.05$). The mean age of starting smoking was 16.95 (SD±4.47) years; for male and female it was 16.83 (SD±4.5) years and 23.66 (SD±10.96) years respectively. The mean number of stick consumed per day was 14.91 (SD±5.82) and it was not varied statistically among male and female. Majority of the smokers (90.6%) smoked cigarette, only a small portion (8.8%) smoked biri and male female distribution was not statistically different ($p > 0.05$). Among the non-daily smokers, past daily smokers were 10.5% and the prevalence was significantly high among male ($p < 0.05$).

Moreover 62.5% and 12.5% non-smokers were exposed to second hand smoke at home or work place respectively and females were significantly more exposed at home than males ($p < 0.05$). Overall prevalence of smokeless tobacco use was 47.5%, the rates were 60.0% and 36.1% among female and male respectively. Smokeless tobacco use was found significantly higher among female ($p < 0.05$). Majority of the current smokeless tobacco users (85.8%) were daily smoker. Most of the smokeless tobacco users use zorda (79.9%), followed by sada pata and guul (10.5% and 9.6%) respectively and male female distribution was not statistically different ($p > 0.05$). Mean frequency of daily smokeless

tobacco use was 8.36 (SD±4.87) and this was not statistically different among male and female ($p > 0.05$).

Based on MET minutes/week, 36.2% of the study subjects fell into low physical activity category (<600 MET minutes/week), 32.8% fell into moderate physical activity category (600–3000 MET minutes/week) and 31% fell into high physical activity group (>3000 MET minutes/week). Prevalence of high physical activity was more in males compared to females and low physical activity was more in females compared to male. This differ statistically ($p < 0.05$). Most of the METs of women were contributed by moderate activities. Almost half of the respondents (46.2%) travelled to and from places by bicycle or by walking and almost all of the respondents (99.1%) did not do any vigorous activity during leisure time. A little more than one fifth of the respondents (21.3%) were engaged in work related vigorous activity, the rates were 37.3% and 3.6% among male and female respectively. Males were engaged in work related vigorous activity significantly more than female ($p < 0.05$). Mean minutes of sedentary activity in a day was 159.76 (SD±71.28), for male it was 163.44 (SD±76.67) and for female, it was 155.73 (SD±64.80) minutes respectively, which did not differ statistically ($p > 0.05$). Almost ninety per cent study subjects (89.6%) were lifetime abstainer of alcohol. Alcohol consumer (drinking wine in last 12 months) was 3.5% and all of them were male. Current drinker (1.7%), daily drinker (0.9%) and binge drinker (0.9%) were also male. Moderate and heavy drinkers were 0.4% each and also were male.

Table 4.1.3: Dietary pattern- fruit, vegetable, oil and salt consumption of the respondents

Variables	Total n=461(100)	Male (n=241)	Female (n=220)	p value
Fruit and vegetable consumption				
Mean (\pm SD) days of fruits intake in a week	2.38 \pm 1.65	2.38 \pm 1.61	2.39 \pm 1.69	0.931
Mean (\pm SD) days of vegetable intake in a week	5.84 \pm 0.88	5.73 \pm 0.920	5.95 \pm 0.837	0.010
Mean (\pm SD) daily fruit intake in servings	1.00 \pm 0.24	1.02 \pm 0.302	0.97 \pm 0.163	0.023
Mean (\pm SD) daily vegetable intake in servings	1.28 \pm 0.45	1.34 \pm 0.474	1.21 \pm 0.411	0.002
Low fruit consumption	461(100)	241(100)	220(100)	
Low vegetable consumption	461(100)	241(100)	220(100)	
Oil consumption				
Types of cooking oil used				
Soyabin oil	371(80.5)	199(82.6)	172(78.2)	0.319
Palm oil	83(18.0)	39(16.2)	44(20.0)	
Mustard oil	5(1.1)	3(1.2)	2(0.9)	
Mean(\pm SD) daily consumption in ml (n=341)	19.12 \pm 5.13	18.87 \pm 5.30	19.27 \pm 5.04	0.494
High oil consumption (n=341, male-122, female-219)	88(25.8)	11(9.0)	77(35.1)	0.000
Salt consumption				
Mean(\pm SD) salt intake per day in gm (n=341, male-122, female-219)	13.26 \pm 3.79	13.32 \pm 4.17	13.24 \pm 3.57	0.855
High salt consumption (n=341, male-122, female-219)	338(99.1)	121(99.2)	217(99.1)	0.929
Extra salt intake(n=461)	448(97.2)	232(96.3)	216(98.2)	0.214

*Values expressed as numbers and percentages in parentheses or $M\pm SD$, as appropriate; M, mean; SD, standard deviation, p value, significance between male and female participants; p value was obtained by χ^2 test for categorical variables & t- test for continuous variable; 0.05 was considered as level of significance.

Table 4.1.3 showed that survey population took fruit on an average 2.38 (SD \pm 1.65) days a week. Vegetables consumption was particularly high. They consumed vegetables in around 5.84 (SD \pm 0.88) days a week. But neither fruit nor vegetable consumption was adequate in quantity. The overall daily per person consumption of fruit was 1.00 (SD \pm 0.24) serving and of vegetables 1.28 (SD \pm 0.45) servings against their minimum daily total requirement of 5 servings. Considering this as minimum recommended

amount, 100% of the study subjects did not consume adequate fruit or vegetables on an average day. All of the 461 study participants could tell about the types of cooking oil used in their home, but only females were able to tell about the quantity of oil and salt they consume in a month. Most of the subjects (80.5%) consume soya bin oil, followed by palm oil (18%), which did not differ statistically among male and female ($p > 0.05$). Of the 241 male, only 122 person could be assessed for their oil and salt consumption and of the 220 female, only one person could not tell about her oil and salt consumption. Average daily oil consumption for male was found 18.87 (SD \pm 5.3) ml and for female it was 19.27 (SD \pm 5.0) ml. This did not vary statistically ($p > 0.05$). For our country, the recommended amount of edible oil to be consumed per day is 5 tsp (25 ml) for men and 4 tsp (20 ml) for women. Only 9% male consume more than recommended level of oil daily, whereas it was 35.2% in case of female. Females took significantly higher amount of oil than male ($p < 0.05$). Total 341 participants could be assessed for their salt consumption. Their mean salt consumption per day was 13.2 (SD \pm 3.79) grams, which was more than double of the recommended level. A total 338 (99.1%) participants consume salt in excess of their normal requirement. Almost all (97.2%) participants took extra salt while taking food and this did not differ statistically among male and female ($p > 0.05$).

Table 4.1.4: Biological risk factors among the respondents

Variables	Total n=461(100)	Male (n=241)	Female(n=220)	<i>p value</i>
Overweight and obesity				
Mean(\pm SD) BMI in kg/m ²	22.37 \pm 2.92	22.2 \pm 2.80	22.55 \pm 3.04	0.202
Category of BMI				
Underweight	32(6.9)	17(7.1)	15(6.8)	0.349
Normal	357(77.4)	192(79.7)	165(75.0)	
Overweight	72(15.6)	32(13.3)	40(18.2)	
Waist circumference				
Mean(\pm SD) waist circumferences in cm	79.83 \pm 7.71	81.36 \pm 8.04	78.16 \pm 6.97	0.000
Category of waist circumferences				
High	130(28.2)	30(12.4)	100(45.5)	0.000
Normal	331(71.8)	211(87.6)	120(54.5)	
Blood pressure				
Mean(\pm SD) systolic blood pressure	115.09 \pm 14.23	117.43 \pm 14.69	112.52 \pm 13.28	0.000
Mean(\pm SD) diastolic blood pressure	73.64 \pm 9.65	75.22 \pm 9.09	71.91 \pm 9.97	0.000
Blood pressure status				
Normal	393(85.2)	201(83.4)	192(87.3)	0.242
Hypertension	68(14.8)	40(16.6)	28(12.7)	0.628
Blood pressure never measured	123(26.7)	62(25.7)	61(27.7)	
Previously diagnosed hypertensive	53(11.5)	28(11.6)	25(11.4)	0.932
Previously diagnosed hypertensive on medication (n=53)	16(30.2)	7(25.0)	9(36.0)	0.384
Antihypertensive medication (allopathic) in last 2 weeks (n=53)	13(24.5)	6(21.4)	7(28.0)	0.579
Herbal or ayurvedic medication (n=53)	3(5.7)	1(3.6)	2(8.0)	0.486
Anti-hypertensive medication among current hypertensive	10(14.7)	6(15.0)	4(14.3)	0.935
Hypertensive unaware about hypertension (n=68, m=40, f=28)	44(64.7)	26(65.0)	18(64.3)	0.952
Hypertension controlled among treatment group (n=16)	6(37.5)	1(14.3)	(55.6)	0.091
Diabetes				
Diabetes	23(5.0)	14(5.8)	9(4.1)	0.397
Blood glucose never measured	404(87.6)	209(86.7)	195(88.6)	0.533
Treatment with insulin	6(1.3)	4(1.7)	2(0.9)	0.478
Treatment with oral anti-diabetic	13(2.8)	8(3.3)	5(2.3)	0.498

**Values expressed as numbers and percentages in parentheses or M±SD, as appropriate; M, mean; SD, standard deviation, p value, significance between male and female participants; p value was obtained by χ^2 test for categorical variables & t- test for continuous variable; 0.05 was considered as level of significance*

Table 4.1.4 showed that mean BMI of respondents was 22.37 (SD±2.92), for male and female it was 22.2 (SD±2.80) and 22.55 (SD±3.04) respectively. More than two third of the population (77.4%) were of normal weight whereas underweight and obesity was 6.9% and 15.6% respectively. Proportion of overweight in women (18.2%) exceeded the proportion of those in men (13.3%). Mean waist circumferences was 79.83 (SD±7.71) cm, for male it was 81.36 (SD±8.04) cm and for female it was 78.16 (SD±6.97) cm. Males mean waist circumference was significantly higher than females ($p < 0.05$). A little more than one fourth of the respondents (28.2%) waist circumference was found high, for male and female, it was 12.4% and 45.5% respectively. The prevalence of high waist circumference was significantly higher in females than in males ($p < 0.05$). Higher prevalence of both central and general obesity in women may predispose them to an increased risk of cardiovascular diseases.

Overall prevalence of hypertension was 14.8% and the rates among male and female were 16.6% and 12.7% respectively. The mean systolic blood pressure of male and female was 117.43 (SD±14.69) mmHg and 112.52 (SD±13.28) mmHg respectively whereas mean diastolic blood pressure of male and female was 75.22 (SD±9.09) mmHg and 71.91 (SD±9.97) mmHg respectively. Both mean systolic and diastolic blood pressure was significantly higher in males than in females ($p < 0.05$). A little less than one third (26.7%) of the study subjects never measured their blood pressure. The prevalence of previously diagnosed hypertension was 11.5%. Among the currently diagnosed hypertensive, 64.7% were unaware of their high blood pressure. Only 14.7% of currently diagnosed hypertensive and 30% of previously diagnosed hypertensive were taking medication. Among treatment group, 81% were taking modern treatment and 19% were taking traditional treatment. A little more than one third (37.5%) of subjects who were on antihypertensive treatment, had blood pressure within normal level. Blood glucose measurement was not included in this study. Documented history of diabetes was sought. Around 87.6% of participants never measured their blood glucose. About 5% of the

people were previously diagnosed to have diabetes (men 5.8% and women 4.1%). Only 1.3% and 2.8% study subjects were treated with insulin and oral anti diabetic agents respectively. Male female distribution was not statistically different ($p > 0.05$).

Table 4.1.5: Family history of coronary heart disease/heart attack and stroke

Variables	Total n=461(100)	Male (n=241)	Female (n=220)	<i>p value</i>
Family history				
Heart attack	2(0.43)	1(0.41)	1(0.45)	
Stroke	7(1.51)	5(2.07)	2(0.90)	0.571

**Values expressed as numbers and percentages; p value, significance between male and female participants; p value was obtained by χ^2 test for categorical variables; 0.05 was considered as level of significance*

Table 4.1.5 showed that almost 98% of the study subjects had no family history of coronary heart disease or stroke and the male female difference was not statistically significant ($p > 0.05$).

4.2 Association between hypertension and various socio-demographic, modifiable and non-modifiable risk factors

The socio-demographic characteristics I studied here are as follows:

1. Age group
2. Education
3. Occupation
4. Family income
5. Sex

Some of these characteristics are also considered as risk factor for CVD like advancing age, sex and low monthly family income.

The risk factors I studied here are as follows:

1. Tobacco use
2. Low fruit and vegetable consumption
3. High salt and oil intake
4. Physical inactivity
5. Harmful alcohol consumption
6. Overweight and obesity
7. High waist circumference
8. Raised blood pressure
9. Diabetes mellitus
10. Advancing age
11. Heredity or family history
12. Gender
13. Low socioeconomic status- low monthly family income.

I could not classify socioeconomic status of study subjects. So instead of low socioeconomic status, I consider low monthly family income as risk factor for CVD.

4.2.1 Association between hypertension and socio-demographic characteristics

Association between hypertension and different socio-demographic characteristics like sex, age group, education, occupation and average monthly family income were sought. For this purpose, education was further categorized into up to primary and primary & above groups. Average monthly family income was categorized into two income groups, one is 10000 taka or less and the other one is above 10000 taka monthly.

Table 4.2.1: Association between hypertension and socio-demographic characteristics

Variables	Hypertension		Total	χ^2	p value
	Present	Absent			
Age group					
35 to <45	16(5.6)	269(94.4)	285(61.8)		
45 to <55	21(18.9)	90(81.1)	111(24.1)		
55 to <65	21(44.7)	26(55.3)	47(10.2)	77.770	0.000
≥65	10(55.6)	8(44.4)	18(3.9)		
Education					
<Primary	50(14.5)	296(85.5)	346(75.1)	0.099	0.753
≥Primary	18(15.7)	97(84.3)	115(24.9)		
Occupation					
House hold work	20(11.0)	161(89.0)	181(39.3)		
Small business	15(15.5)	82(84.5)	97(21.0)		
Day labourer	6(10.9)	49(89.1)	55(11.9)		
Driver	3(10.3)	26(89.7)	29(6.3)	35.743	0.001
Rickshaw puller	2(11.1)	16(88.9)	18(3.9)		
Private employee	4(23.5)	13(76.5)	17(3.7)		
Family income					
≤10000	8(4.7)	161(95.3)	169(36.7)		
>10000	60(20.5)	232(79.5)	292(63.3)	21.289	0.000
Sex					
Male	40(16.6)	201(83.4)	241(52.3)		
Female	28(12.7)	192(87.3)	220(47.7)	1.370	0.242

Table 4.2.1 showed statistically significant association between hypertension and age group ($p < 0.001$), occupation ($p < 0.01$) and monthly family income ($p < 0.001$). No association was found for education and gender with hypertension.

4.2.2 Association between hypertension and modifiable risk factors

Association between hypertension and modifiable risk factors like tobacco use, oil consumption, physical activity, nutritional status based on BMI, central obesity based on waist circumference and previously diagnosed diabetes were sought. Low fruit and vegetable consumption was not studied because all of the respondents fell into low fruit and vegetable consumption group. Salt consumption could not be analyzed because 99.1% of the respondents who could tell about their salt consumption were high salt consumer. Alcohol consumption was also omitted because a very small percentage of the respondents were alcohol drinker.

Table 4.2.2: Association between hypertension and modifiable risk factors

Variables	Hypertension		Total	χ^2	p value
	Present	Absent			
Smoking tobacco use	24(13.3)	157(86.7)	181(39.3)	0.527	0.468
smokeless tobacco use	42(19.2)	177(80.8)	219(47.5)	6.504	0.011
Second hand smoking-home	30(17.1)	145(82.9)	175(38.0)	1.284	0.257
Second hand smoking - workplace	5(14.3)	30(85.7)	35(7.6)	0.007	0.936
High oil consumption in male (n=122)	2(18.2)	9(81.8)	11(9.0)	0.004	0.952
High oil consumption in female (n=219)	10(13)	67(87)	77(35.2)	0.004	0.948
Physical activity					
High	15(10.5%)	128(89.5%)	143(31.0%)	9.682	0.008
Moderate	17(11.3%)	134(88.7)	151(32.8)		
Low	36(21.6)	131(78.4)	167(36.2)		
Nutritional status- BMI					
Underweight	2(6.3)	30(93.7)	32(6.9)	24.142	0.000
Normal	42(11.8)	315(88.2)	357(77.4)		
Overweight and obese	24(33.3)	48(66.7)	72(15.6)		
High waist circumference- male (n=241)	12(40.0)	18(60.0)	30(12.4)	13.557	0.000
High waist circumference- female (n=220)	19(19.0)	81(81.0)	100(45.5)	2.873	0.090
Diabetes	11(47.8)	12(52.2)	23(5.0)	21.061	0.000

Statistical significant association was found between hypertension and BMI ($p < 0.001$), physical activity ($p < 0.005$), smokeless tobacco use ($p < 0.01$), diabetes ($p < 0.001$) and waist circumference in male ($p < 0.001$).

4.2.3 Association between hypertension and non-modifiable risk factors

Non-modifiable risk factors like advancing age and sex were discussed in socio-demographic section. Only family history of heart attack or stroke is analyzed here.

Table 4.2.3: Association between hypertension and family history of heart attack or stroke

Variable	Hypertension		Total	χ^2	p value
	Present	Absent			
Family history (+)	2(22.2)	7(77.8)	9(2.0)	0.408	0.523
Family history (-)	66(14.6)	386(85.4)	452(98.0)		

The table 4.2.3 showed no association between hypertension and family history of heart attack or stroke ($p > 0.05$).

4.3 Effect of various modifiable and non-modifiable risk factors on hypertension

Modifiable risk factors I tested include current smoker, current smokeless tobacco user, low physical activity, overweight and obesity, high waist circumference, low monthly family income and previously diagnosed diabetes. Low fruit and vegetable consumption was not tested because all of the respondents fell into low fruit and vegetable consumption group. Salt and oil consumption was not included because a large portion of the respondents did not give proper history regarding salt and oil consumption. Alcohol consumption was also omitted because a very small percentage of the respondents were alcohol drinker. Non modifiable risk factors include gender, advancing age, and family history of coronary heart disease/stroke. Family history was not tested because a very negligible portion of the respondents had a positive family history. For convenience of the study, age was further categorized into above 55 years and 55 years or less. Weight was categorized again into normal to underweight and overweight & obese group. Physical activity was further categorized into low physical activity and physically active group. Both high and moderate physical activity were combined to categorize as physically active group,

Table 4.3.1: Effect of various modifiable and non-modifiable risk factors on hypertension

Variables	Crude OR	95% CI		Adjusted OR	95% CI	
		Lower	Upper		Lower	Upper
Sex	1.365	.810	2.300	2.761	1.136	6.712
Age group	8.847	4.890	16.004	6.273	3.112	12.644
Monthly family income	5.205	2.423	11.181	3.834	1.668	8.815
Current smoker	.820	.479	1.402	.593	.256	1.374
Current smokeless tobacco user	1.971	1.163	3.342	2.598	1.316	5.126
Physical activity	2.250	1.337	3.786	1.895	1.007	3.567
Overweight and obesity	3.920	2.191	7.015	2.284	1.048	4.978
High waist circumference	2.488	1.466	4.222	2.170	1.012	4.654
Diabetes	6.127	2.582	14.540	2.004	.659	6.095

**Reference category was female sex, age less than 55 years, monthly family income 10000 taka or less, non smoker, not smokeless tobacco user, physically active, normal or underweight, normal waist circumference and non diabetic.*

The table 4.3.1 showed that after adjusting all other effects, risk of hypertension was found almost 3 times higher in male compared to female (OR 2.761, 95%CI 1.136-6.712). Risk of hypertension was more than six times higher in higher age group compared to lower age group (OR 6.273, 95%CI 3.112- 12.644). High income group was found to have almost 4 times higher risk of hypertension compared to low income group (OR 3.834, 95%CI 1.668-8.815). Current smokeless tobacco users were also found to have more than 2.5 times higher risk of hypertension than non smokers (OR 2.598, 95%CI 1.316-5.126). Risk of hypertension was found to be almost 2 times higher among the people with low physical activity compared to high activity group (OR 1.895, 95%CI 1.007-3.567). Risk of hypertension was also found to be more than 2 times higher in overweight-obese (OR 2.284, 95% CI 1.048-4.978) and high waist circumference (OR 2.170, 95% CI 1.012-4.654) group compared to normal to underweight and normal waist circumference group respectively. No significant effect was found for current smoking status and diabetes on hypertension.

4.4 Effect of non-modifiable risk factors on various modifiable risk factors

Non modifiable risk factors include gender, advancing age, and family history of coronary heart disease/stroke. Family history was not tested because a very negligible portion of the respondents had a positive family history. For convenience of the study, age was further categorized into above 55 years and 55 years or less.

Table 4.4.1: Effect of sex and age on current smoking status

Variables	Crude OR	95% CI		Adjusted OR	95% CI	
		Lower	Upper		Lower	Upper
Sex	149.344	53.343	418.114	166.284	58.821	470.081
Age group	1.495	.882	2.534	.533	.275	1.032

**Reference category was female sex and age less than 55 year.*

Table 4.4.1 showed that after adjustment, risk of smoking was 166 times more in male compared to female (OR 166.28, 95%CI 58.82- 470.08). Age had no effect on smoking.

Table 4.4.2: Effect of sex and age on current smokeless tobacco user

Variables	Crude OR	95% CI		Adjusted OR	95% CI	
		Lower	Upper		Lower	Upper
Sex	.377	.258	.549	.362	.246	.532
Age group	1.009	.597	1.706	1.341	.773	2.325

**Reference category was female sex and age less than 55 year.*

Table 4.4.2 showed that after adjustment, risk of smokeless tobacco use was 74% less in male compared to female (OR 0.362, 95%CI 0.246-0.532). Age had no effect on smokeless tobacco use.

Table 4.4.3: Effect of sex and age on physical activity

Variables	Crude OR	95% CI		Adjusted OR	95% CI	
		Lower	Upper		Lower	Upper
Sex	0.731	0.500	1.071	.631	.424	.939
Age group	2.333	1.373	3.966	2.670	1.544	4.619

**Reference category was female sex and age less than 55 year.*

Table 4.4.3 showed that after adjustment, risk of low physical activity was almost 37% less in male compared to female (OR 0.631, 95%CI 0.424- 0.939) and more than 2.5 times higher in 55 years or more age group compared to less than 55 years age group (OR 2.670, 95%CI 1.544- 4.619).

Table 4.4.4: Effect of sex and age on overweight and obesity

Variables	Crude OR	95% CI		Adjusted OR	95% CI	
		Lower	Upper		Lower	Upper
Sex	.689	.415	1.143	.641	.381	1.078
Age group	1.550	.792	3.033	1.620	.814	3.221

**Reference category was female sex and age less than 55 year.*

Table 4.4.4 showed that there was no significant effect of sex and age on overweight and obesity.

Table 4.4.5: Effect of sex and age on waist circumference

Variables	Crude OR	95% CI		Adjusted OR	95% CI	
		Lower	Upper		Lower	Upper
Sex	.171	.107	.272	.148	.091	.243
Age group	1.156	.653	2.045	2.168	1.127	4.173

**Reference category was female sex and age less than 55 year.*

Table 4.4.5 showed that after adjustment, risk of high waist circumference was almost 85% less in male compared to female (OR 0.148, 95%CI 0.091- 0.243) and more than 2 times higher in 55 years or more age group compared to less than 55 years age group (OR 2.168, 95%CI 1.127- 4.173).

Table 4.4.6: Effect of sex and age on diabetes

Variables	Crude OR	95% CI		Adjusted OR	95% CI	
		Lower	Upper		Lower	Upper
Sex	1.446	.613	3.410	.919	.366	2.310
Age group	7.925	3.330	18.860	8.109	3.283	20.034

**Reference category was female sex and age less than 55 year.*

Table 4.4.6 showed that after adjustment, risk of diabetes was more than eight times higher in 55 years or more age group compared to less than 55 years age group (OR 8.11, 95%CI 3.28- 20.03). Sex had no effect on diabetes.

CHAPTER 5
DISCUSSION

Chapter 5

Discussion

To the best of my knowledge, this is the first CVD risk factor survey among urban slum population in Bangladesh. The study subjects, mostly rural migrants who had come to Dhaka for economic reasons and residing in the slums showed a marked prevalence of hypertension and unawareness about hypertension status, very high prevalence of smoking tobacco use in male and smokeless tobacco use in female, sedentary life style and low physical activity with very minimum exercise, very much inadequate fruit and vegetable consumption with high salt intake with invariable use of extra salt in diet and higher prevalence of high waist circumference among female.

Cardiovascular disease (CVD) is a rising health burden among the world's poor with hypertension as the main risk factor (102). Slums are typically characterized by poor living conditions with limited access to quality healthcare. The prevalence of risk factors for CVD is high in slums and the associated psychosocial burden of insecurity, violence and stress may cause an increased risk of CVD. Therefore a large part of the CVD burden is on the urban poor, who may not have the financial resources or adequate health literacy. In India, people in slums are generally at a higher risk for CHD than Indians living in more affluent areas mostly because of the higher prevalence of major CHD risk factors such as uncontrolled hypertension and tobacco use amongst them (103).

In the current study, the prevalence of hypertension was found to be 14.8% and the rates among male and female were 16.6% and 12.7% respectively. Among the hypertensive, 64.7% were unaware of their diagnosis, only 14.7% of currently diagnosed hypertensive and 30.2% of previously diagnosed hypertensive were taking medication and 37.5% of treatment group had controlled blood pressure. Mean systolic blood pressure was 115.09 (SD \pm 14.23) and mean diastolic blood pressure was 73.64 (SD \pm 9.65). Both systolic and diastolic blood pressure was high in males compared to females. A cross-sectional, population-based study was conducted to assess the prevalence, awareness, treatment,

and control of hypertension among people aged 20-59 years from 120 clusters spread across Delhi. According to this study, the prevalence of hypertension among slum dwellers was 23%. Among the hypertensive, 55% were unaware of their diagnosis; about 36% were taking treatment and only 9.4% had controlled blood pressure (BP) (103). Studies conducted in slums of Faridabad, Delhi and urban Chennai in India among lower socioeconomic groups reported a prevalence of 17.2, 12 and 8.4 % respectively (104–106). Two studies in Kenya have reported on hypertension among poor urban slum residents. Of these one was among Kibera slum residents and reported an unadjusted hypertension prevalence of 13%, the rates were 17.8% and 11.1% among males and females respectively. The mean blood pressure was 122/71mmHg. Among the hypertensive, 87% were unaware of their diagnosis and only 10% of them were taking treatment (107). The other study, a population based survey in an urban slum in Nairobi, Kenya showed prevalence of hypertension was 22.8%, among them 80% were unaware of their hypertensive status (108). Findings of these studies were consistent with the findings of the current study. In another population based study on urban slum dwellers done in Patna, India showed prevalence of hypertension were 16.36 percent (males 18.79 %, females 14.48%. (109). This study finding is compatible with the current study. A descriptive study among slum people in Nairobi, Kenya showed the mean systolic and diastolic blood pressure was 149.2 and 95.2 mmHg respectively. Among hypertensive, only 6.8% were aware about their hypertension status (102). The mean systolic and diastolic blood pressure was much less in the current study. Dissimilar result also found regarding awareness about hypertension status.

Moreover, statistical significant association was found between hypertension and advancing age, higher monthly family income, increased BMI, low physical activity, smokeless tobacco use, diabetes and increased waist circumference in male sex in the current study. Study conducted in Patna by Singh et al showed statistical significant association ($p < 0.05$) of hypertension with BMI, waist circumference, tobacco use, alcohol consumption and physical inactivity (109). Kibera slum study showed significant association of hypertension with male gender, high body mass index and increasing age

(107). Joshi MD et al showed that older age, higher general and central obesity were independently associated with hypertension (108).

Logistic regression in current study showed increased risk of hypertension among male sex, higher age group, high income group, smokeless tobacco use, low physical activity, high BMI and high waist circumference. Multivariate logistic regression by Singh et al revealed that high BMI and alcohol were associated with hypertension (109). Another slum study in India showed increased risk of hypertension with higher age group. Findings of these studies are consistent with current study (110).

The current study showed a high prevalence of smoking tobacco use in male (39.3% in both sex with 73.4% among male sex) and a strikingly high prevalence of smokeless tobacco use (60%) in female. Singh et al showed tobacco users were 12.54 percent (109). Joshi MD et al showed current smokers were 10% (108). A study in Bihar, India showed 22.75 percent prevalence of smoking in both sexes and all social classes (111) The prevalence of smoking is much higher in current study. Another study in North India showed higher prevalence of tobacco use among slum dwellers (men 48.3%, women 11.9%) than non-slum dwellers (men 35.2%, women 3.5%). (11) This study finding is compatible with the current study. Several cohort and case-control studies have also reported a significant positive association between smokeless tobacco use and CHD (112–114). This study also showed smokeless tobacco use as significant predictor of hypertension (OR2.598, 95%CI 1.316-5.126).

Though majority of the study subjects (77.4%) in the current study had normal BMI, the overall prevalence of overweight and obesity was 15.6%. Female showed a higher prevalence of overweight compared to male (18.2% versus 13.3%). Also females had much higher prevalence of central obesity (45.5%). Study done in Patna, India showed higher prevalence of high body mass index (BMI) and waist circumference (WC) than the current study; 31.94 percent (males 31.83%, females 32.03%) and 50.45 percent (males 39.1%, females 59.17%) respectively (109). In urban Chennai, Mohan et al reported a 33 % prevalence of overweight and obesity in low income group and the

prevalence increased with increase in income (106). Misra et al and ICMR Task force study reported a prevalence of overweight of 25 percent and 20 percent respectively in Delhi slums (105,115). The study conducted in Patna, India showed higher prevalence of central obesity in females as compared to males (109). A study in Kenya showed 58.3% of women were either obese or over-weight and 41.5% females had high waist circumference (107). The current study also showed similar result (45.5% in females versus 12.4% in males). The study done by Joshi MD et al showed majority of males had normal BMI and waist circumference, whereas a third of females were obese or overweight and 40% had central obesity (108). Findings were mostly consistent with the current study. An observational study conducted on women in a slum in Peru showed prevalence of central obesity and BMI obesity were 90% and 30% respectively (116). The present study showed a lesser prevalence of central and BMI obesity. A descriptive study among slum people in Nairobi, Kenya showed a higher BMI in 63% of women and higher waist circumference more in men (102). The present study showed lesser prevalence of higher BMI in both sexes and increased waist circumference more in women.

Singh et al showed significant association of hypertension with both central and general obesity ($p < 0.05$). Multivariate logistic regression revealed that high BMI were associated with hypertension (109). BMI levels of >25 was significantly associated with hypertension in the current study; similar results were observed in Indians living in Mauritius who had increased rates of HTN at these BMI levels (117). Chirinos et al. of Pennsylvania school of Medicine reports among patients with hypertension increasing BMI was a significant predictor of hypertension (OR1.04,95%CI,1.02-1.06) (118). The current study also showed similar result (OR2.28, 95%CI 1.05-4.98). Study in Kenya also showed independent association of hypertension with higher general and central obesity (108).

The present study showed prevalence of low physical activity, alcohol consumption and diabetes as 36.2%, 3.5% and 5% respectively. Prevalence of low physical activity was more in females than males (40% versus 33%). The urban slum study in Nairobi, Kenya

showed prevalence low physical activity, harmful alcohol drinking and diabetes were 20%, 52% and 5% respectively (108). Alcohol consumption was much less in the current study. This is because our society is much conservative and people are mostly followers of religious law. Almost all of the alcohol consumers were found hesitant to tell about their alcohol drinking.

The increased prevalence of CHD and its risk factors among slum dwellers is of major concern since this group is large in number, and they have poor access to acute care management and long-term secondary prevention practices. This is due to the inadequacy of the urban public health delivery system to access these areas of very high population density. In spite of better health care infrastructure, primary healthcare facilities have not grown in proportion to the explosive growth of urban population, in particular the slum population. The social exclusion and lack of information and assistance at the secondary and tertiary hospitals make them unfamiliar with the modern hospital environment making them less likely to use these facilities. The lack of economic resources also restricts their access to the available private facilities. Moreover, a general lack of understanding about the causes of CHD and social stigma among this group results in a large proportion of those with illness remains undiagnosed and very few undertaking preventive medical treatment. Similarly, as they have poor compliance and adherence to pharmaceutical treatments, they are also more exposed to future CHD events. A continued neglect of the healthcare of people in overcrowded urban slums will inevitably lead to greater expenditure and diversion of healthcare resources in the long-term, having to manage end stage complications of major risk factors for CHD. Research has confirmed that CHD risk factor occurs frequently in cluster in hypertensive individuals and that the risk of developing CHD is determined by the cumulative effect of multiple concomitant CHD risk factors (119). Therefore, targeting preventive intervention at these people is fundamental to an effective CHD prevention approach.

CHAPTER 6

Conclusion, Limitation and Recommendations

Chapter 6

Conclusion, Limitation and Recommendations of the Study

6.1 Conclusion

The study population showed high prevalence of hypertension and other cardiovascular disease risk factors like both smoking and smokeless tobacco use, low physical activity with very minimum exercise and a considerable amount of time of sedentary activity, very minimum fruit and vegetable intake with very high salt intake. Two important findings were very high smokeless tobacco use and high waist circumference among female. Furthermore, the vast majority of hypertensive patients were unaware about their hypertension status and very small portion of them were on medication. Effectiveness of treatment was also under question, because hypertension was not controlled in majority of patients. These findings points to the need for greater awareness of hypertension and other cardiovascular disease risk factors in this population. Preventive efforts in this population targeting behavioural changes in lifestyle to curtail these risk factors urgently required. Reducing these risk factors will not only have an effect on cardiovascular diseases, but will also have wide-reaching beneficial effects for other chronic non-communicable diseases.

6.2 Limitations

Limitations of studies are very common in social work. This study also has some limitations. First of all, due to budget constraints, I only did the first two steps of WHO steps approach- questionnaire and physical measurement. The third step- biochemical measurement of blood glucose and lipid profile could not be done. Secondly, self reporting and recall bias might have affected the validity of response to questions wealth, education, occupation, smoking, alcohol misuse, diet and physical activity. Self-reporting can lead to inaccurate reporting due to lack of awareness, misinterpretation of questions, or concern for judgement and affect study conclusions. Recall bias in dietary assessment is very much common. In this study only females were able to tell about their salt and oil consumption. Thirdly, full dietary assessment was not done. Finally, in clinical practice, a diagnosis of hypertension requires multiple measurements on several occasions. I took a single occasion measurement and therefore, the prevalence of hypertension found in this survey may represent an over or underestimation.

6.3 Recommendations

On the basis of key findings, the recommendations would be followings:

- Further large scale study is recommended including other slums which will represent all slum population.
- Levels of health literacy have to increase in order for people to change their lifestyle, get diagnosed or comply with therapy. Health education program focusing on smoking, physical activity, dietary pattern and importance of blood pressure management may be introduced.
- Follow up study is necessary to determine the prediction capability of the tool.

CHAPTER 7
REFERENCES

Chapter 7

References

1. WHO Cardiovascular diseases fact sheet [Internet]. 2015. Available from: <http://www.who.int/mediacentre/factsheets/fs317/en/>
2. Mathers CD, Loncar D. Projections of global mortality and burden of disease from 2002 to 2030. *PLoS Med.* 2006;3(11):2011–30.
3. Global status report on noncommunicable diseases. WHO. 2010;ISBN: 978 92 4 156422 9.
4. Ogunmola OJ, Olaiḡa AO, Oladapo OO, Babatunde O a. Prevalence of cardiovascular risk factors among adults without obvious cardiovascular disease in a rural community in Ekiti State, Southwest Nigeria. *BMC Cardiovasc Disord* [Internet]. *BMC Cardiovascular Disorders*; 2013;13(1):89. Available from: <http://www.pubmedcentral.nih.gov/articlerender.fcgi?artid=4016363&tool=pmcentrez&rendertype=abstract>
5. World Health Organization (WHO). who ncd country profile bangladesh 2008 [Internet]. 2008. Available from: www.who.int/nmh/countries/bgd_en.pdf
6. World Health Organization. Non-Communicable Disease Risk Factor Survey Bangladesh 2010 [Internet]. 2011. Available from: http://www.searo.who.int/bangladesh/publications/ncd_risk_ractor_survey_report_ban_2010.pdf
7. Saquib N, Saquib J, Ahmed T, Khanam M, Cullen MR. Cardiovascular diseases and Type 2 Diabetes in Bangladesh: A systematic review and meta-analysis of studies between 1995 and 2010. *BMC Public Health* [Internet]. *BMC Public Health*; 2012;12(1):434. Available from: *BMC Public Health*
8. Muhit MA, Rahman MO, Raihan SZ, Asaduzzaman M, Akbar MA, Sharmin N, et al. Cardiovascular disease prevalence and prescription patterns at a tertiary level hospital in Bangladesh. *J Appl Pharm Sci.* 2012;2(03):80–4.

9. Robert Beaglehole AI and TP. The World health report : 2003 : shaping the future. [Internet]. World Health Organization. 2003. Available from: www.who.int/whr/2003/en/whr03_en.pdf
10. JI C, Dphil M. Mortality by cause for eight regions of the world : Global Burden of Disease Study. *Lancet*. 1997;349(9061):1269–76.
11. Gu D, Gupta A, Muntner P, Hu S, Duan X, Chen J, et al. Prevalence of cardiovascular disease risk factor clustering among the adult population of China: Results from the International Collaborative Study of Cardiovascular Disease in Asia (InterAsia). *Circulation*. 2005;112(5):658–65.
12. Batis C, Gordon-larsen P, Cole SR, Du S, Zhang B, Popkin B. Hypertension Among Chinese Adults. 2013;24(3):1–9.
13. Gu DF1, Jiang H, Wu XG, Reynolds K, Gan WQ, Liu DH, Su SY, Duan XF, Huang GY WP. Prevalence, awareness, treatment and control of hypertension in Chinese adults. *Zhonghua Yu Fang Yi Xue Za Zhi* 2003 Mar;37(2)84-9. 2003;37(2):84–9.
14. He J1, Gu D, Reynolds K, Wu X, Muntner P, Zhao J, Chen J, Liu D, Mo J WPICG. Serum total and lipoprotein cholesterol levels and awareness, treatment, and control of hypercholesterolemia in China.No Title. *Circ* 2004 Jul 27;110(4)405-11. 2004;110(4):405–11.
15. Health NYSD of. The burden of Cardiovascular disease in New York: mortality, prevalence, risk factors, costs, and selected population [Internet]. 1999. Available from: https://www.health.ny.gov/diseases/.../heart.../burden_of_cvd_in_nys.pdf
16. Steps WHO, Manual S. WHO STEPwise approach to non-communicable disease risk factor surveillance (STEPS) [Internet]. 2008. Available from: <http://www.who.int/chp/steps/en/>
17. Guindon GE, Guindon GE, Boisclair D, Boisclair D. Past, Current and Future Trends in Tobacco Use. *Population* (English Edition). 2003.
18. Robert AH GW. Morbidity and Mortality Weekly Report (MMWR): Cardiovascular Disease Risk Factors and Preventive Practices Among Adults -- United States, 1994 A Behavioral Risk Factor Atlas [Internet]. National Center for Chronic Disease Prevention and Health Promotion. Available from: <http://www.cdc.gov/mmwr/preview/mmwrhtml/00055888.htm>

19. Guilbert JJ. The world health report 2002 - reducing risks, promoting healthy life. Educ Health (Abingdon). 2003;16(2):230.
20. World Health Organization. Preventing chronic diseases : a vital investment : WHO global report. [Internet]. World Health Organization. 2005. Available from: http://www.who.int/chp/chronic_disease_report/full_report.pdf
21. World Heart Federation. Cardiovascular disease Risk factors. Cardiol J [Internet]. 2012;(April 2012):6–9. Available from: <http://www.world-heart-federation.org/press/fact-sheets/cardiovascular-disease-risk-factors/>
22. Karvosen MJ, Barry a. J. Physical Activity and the Heart. Am J Med Sci. 1967;254(1):126.
23. O'Connor GT, Buring JE, Yusuf S, Goldhaber SZ, Olmstead EM, Paffenbarger RS, et al. An overview of randomized trials of rehabilitation with exercise after myocardial infarction. Circulation. 1989;80(2):234–44.
24. U.S. Department of Health, Human Services C for DC and P. Physical Activity and Health: A Report of the Surgeon General. Atlanta [Internet]. U.S. Department of Health and Human Services, Centers for Disease Control and Prevention, National Center for Chronic Disease Prevention and Health Promotion. 1996. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/20467523>
25. World Health Organization (WHO). Global status report on alcohol 2004. 2004;1–94. Available from: <http://scholar.google.com/scholar?hl=en&btnG=Search&q=intitle:Global+Status+Report+on+Alcohol+2004#0>
26. Koehlmoos TP, Uddin MJ, Ashraf A, Rashid M. Homeless in Dhaka: Violence, sexual harassment, and drug-abuse. J Heal Popul Nutr. 2009;27(4):452–61.
27. Foster WR, Burton BT eds. National Institutes of Health Consensus Development Panel on the health implications of obesity. Ann Intern Med. 1985;103(6):1073–7.
28. International Diabetes Federation (IDF). 5th edition Diabetes Atlas: One adult in ten will have diabetes by 2030 [Internet]. 2011. Available from: <http://www.idf.org/media-events/press-releases/2011/diabetes-atlas-5th-edition>
29. Kannel WB, McGee DL. Diabetes and cardiovascular risk factors: The Framingham study (Article). Circulation [Internet]. 1979;59(1):8–13. Available from:

- <http://www.scopus.com/record/display.url?eid=2-s2.0-0018341014&origin=inward&txGid=8571E4C65D01DE91BAAF5D63A6B89E73.ZmAySxCHIBxxTXbnsoc5w:2>
30. Engelgau MM, Narayan KM, Herman WH. Screening for type 2 diabetes. *Diabetes Care*. 2000;23(10):1563–80.
 31. Akter S, Rahman MM, Abe SK, Sultana P. Prevalence of diabetes and prediabetes and their risk factors among Bangladeshi adults: a nationwide survey. *Bull World Health Organ* [Internet]. 2014;92(3):204–13, 213A. Available from: <http://www.pubmedcentral.nih.gov/articlerender.fcgi?artid=3949596&tool=pmcentrez&rendertype=abstract>
 32. Franceschini G. Epidemiologic evidence for high-density lipoprotein cholesterol as a risk factor for coronary artery disease. *Am J Cardiol*. 2001;88((12A)):9N – 13N.
 33. Cullen P. Evidence that triglycerides are an independent coronary heart disease risk factor. *Am J Cardiol*. 2000;86(9):943–9.
 34. WHO. The Atlas of Heart Disease and Stroke, WHO.Risk factor: socioeconomic status [Internet]. 2011. Available from: http://www.who.int/cardiovascular_diseases/en/cvd_atlas_11_socioeconomics.pdf?ua=1
 35. WHO. The Atlas of Heart Disease and Stroke, WHO [Internet]. Available from: http://www.who.int/cardiovascular_diseases/resources/atlas/en/
 36. World economic forum report on the total global costs of non-communicable diseases (NCDs). [Internet]. Available from: <http://www.weforum.org/news/non-communicable-diseases-cost-47-trillion-2030-new-study>.
 37. icddr b. Manoshi Community Health Solutions in Bangladesh (Impact Evaluation Surveys in Dhaka Urban Slums 2007, 2009 and 2011). 2011.
 38. WHO. World health statistics 2009. Geneva: World Health Organization; 2009e.
 39. Kelly VF and BB. Promoting Cardiovascular Health in the Developing World. INSTITUTE OF MEDICINE OF THE NATIONAL ACADEMIES; 2009. 49 p.
 40. Beaglehole R BR. Global public health: A scorecard. *Lancet*. 372(9654):1988–96.
 41. World Health Organization N. The Global Burden of Disease: 2004 update. Update [Internet]. 2008;2010:146. Available from: http://www.who.int/healthinfo/global_burden_disease/2004_report_update/en/index.html

42. Hardoon SL, Whincup PH, Lennon LT, Wannamethee SG, Capewell S MR. How much of the recent decline in the incidence of myocardial infarction in British men can be explained by changes in cardiovascular risk factors? Evidence from a prospective population-based study. *Circ* 2008. 2008;117(5):598–604.
43. Mirzaei M, Truswell AS, Taylor R LS. Coronary heart disease epidemics: Not all the same. *Heart*. 2009;95(9):740–6.
44. Xavier D, Pais P, Devereaux PJ, Xie C, Prabhakaran D, Reddy KS GR. Treatment and outcomes of acute coronary syndromes in India (CREATE): a prospective analysis of registry data. *Lancet*. 2008;371:1435–42.
45. Gupta R, Joshi P, Mohan V, Reddy KS YS. Epidemiology and causation of coronary heart disease and stroke in India. *Heart*. 2008;94:16–26.
46. Ebrahim S, Kinra S, Bowen L, Andersen E, Ben-Shlomo Y, Lyngdoh T et al. The effect of rural-to-urban migration on obesity and diabetes in india: A cross-sectional study. *PLoS Med*. 2010;7.
47. Deb S DA. A Study on Risk Factors of Cardiovascular Diseases in an Urban Health Center of Kolkata. *Indian J Community Med*. 2008;33(4):271–5.
48. McKeigue PM, Ferrie JE, Pierpont T MM. Association of early-onset coronary heart disease in South Asian men with glucose intolerance and hyperinsulinemia. *Circulation*. 1993;87:152–61.
49. National Heart Foundation Hospital & Research Institute. No Titl [Internet]. Available from: www.nhf.org.bd
50. L. D. Preventing Coronary Heart Disease Risk of Slum Dwelling Residents in India. *J Fam Med Prim Care*. 3(1):58–62.
51. Pencina MJ, D'Agostino RB, Larson MG, Massaro JM VR. Predicting the 30-year risk of cardiovascular disease: The Framingham Heart Study. *Circulation*. 2009;119(24):3078–84.
52. Hunink MG, Goldman L, Tosteson AN, Mittleman MA, Goldman PA, Williams LW, Tsevat J WM. The recent decline in mortality from coronary heart disease, 1980–1990. The effect of secular trends in risk factors and treatment. *J Am Med Assoc*. 1997;277(7):535–42.
53. Yach D, Leeder SR, Bell J KB. Global chronic diseases. *Science*. 2005;307:317.

54. Lopez AD, Mathers CD, Eszati M, Jamison DT MCWB. Global burden of disease and risk factors. Washington, DC: World Bank; 2006.
55. Ezzati M, Henley SJ, Thun MJ LA. Role of smoking in global and regional cardiovascular mortality. *Circulation*. 2005;112(4):489–97.
56. Gritz ER, Vidrine DJ CFM. Smoking cessation. A critical component of medical management in chronic disease populations. *Am J Prev Med*. 2007;33:S414–22.
57. Pilote L, Dasgupta K, Guru V, Humphries KH MJ. A comprehensive view of sex-specific issues related to cardiovascular disease. *Can Med Assoc Journal*. 2007;176(6):1310.
58. Mendis S, Puska P NB editors. *Global Atlas on Cardiovascular Disease Prevention and Control*. Geneva; 2011.
59. He FJ MG. A comprehensive review on salt and health and current experience of worldwide salt reduction programmes. *J Hum Hypertens*. 2009;23(6):363–84.
60. Karppanen H ME. Sodium intake and hypertension. *Prog Cardiovasc Dis*. 2006;49:59–75.
61. Lucas DL, Brown RA, Wassef M GT. Alcohol and the cardiovascular system research challenges and opportunities. *J Am Coll Cardiol*. 2005;45(12):1916–24.
62. Foster RK MH. Alcohol consumption in the new millennium—weighing up the risks and benefits for our health. *Nutr Bull*. 2006;31(4):286–331.
63. Services USD of H and H. *Physical activity guidelines advisory committee report*. Washington, DC; 2008.
64. Guthold R, Ono T, Strong KL, Chatterji S MA. Worldwide variability in physical inactivity—a 51-country survey. *Am J Prev Med*. 2008;34(6):486–94.
65. joint WHO/FAO expert committee. *Joint WHO/FAO Expert Consultation on Diet Nutrition and the Prevention of Chronic Diseases*. Geneva; 2003.
66. Chatterjee S, Allan R RR. Discussion paper no. 07.05: Food consumption and calorie intake in contemporary India. Palmerston North, New Zealand: Massey University Department of Applied and International Economics; 2007.
67. Wang L, Qi X, Chen C, Ma J, Li L, Rao K, Kong L, Ma G, Xiang H PJ. Survey and report on the status of nutrition and health of the Chinese people: A comprehensive report for 2002. Beijing; 2005.

68. Barquera S, Hernandez-Barrera L, Tolentino ML, Espinosa J, Shu WN, Rivera JA PB. Energy intake from beverages is increasing among Mexican adolescents and adults. *J Nutr.* 2008;138(12):2454–61.
69. Henderson L, Gregory J, Irving K SG. National diet & nutrition survey: Adults aged 19 to 64: Energy, protein, carbohydrate, fat, and alcohol intake. London; 2003.
70. Lawes CMM, Vander Hoorn S RA. International Society of Hypertension. Global burden of blood-pressure-related disease,2001 . *Lancet.* 2008;371(9623):1513–8.
71. Gaziano TA, Reddy KS, Paccaud F, Horton S C V. Cardiovascular disease. Disease control priorities in developing countries. 2nd ed. New York: Oxford University Press; 2006. 645–662. p.
72. Mayosi BM, Flisher AJ, Lalloo UG, Sitas F, Tollman SM BD. The burden of non-communicable diseases in South Africa. *Lancet.* 2009;374(9693):934–47.
73. Kannel WB, Castelli WP, Gordon T MP. Serum cholesterol, lipoproteins, and the risk of coronary heart disease. The Framingham Study. *Ann Intern Med.* 1971;74(1):1–12.
74. Jr. GA. Evolving concepts of dyslipidemia, atherosclerosis, and cardiovascular disease: *J Am Coll Cardiol.* 2005;46(7):1219–24.
75. Downs JR, Clearfield M, Weis S, Whitney E SD. Primary prevention of acute coronary events with lovastatin in men and women with average cholesterol levels: Results of AFCAPS/TEXCAPS. Air Force/Texas Coronary Atherosclerosis Prevention Study. *J Am Med Assoc.* 1998;279(20):1615–22.
76. Collins R, Armitage J, Parish S, Sleight P PR. Effects of cholesterol-lowering with simvastatin on stroke and other major vascular events in 20536 people with cerebrovascular disease or other high-risk conditions. *Lancet.* 2004;363(9411):757–67.
77. Yusuf PS, Hawken S, Ounpuu S, Dans T AA. Effect of potentially modifiable risk factors associated with myocardial infarction in 52 countries (the INTERHEART study): Case-control study. *Lancet.* 2004;364(9438):937–52.
78. Asia Pacific Cohort Studies Collaboration. The effects of diabetes on the risks of major cardiovascular diseases and death in the Asia Pacific region. *Diabetes Care.* 2003;26(2):360–6.

79. Kengne AP, Nakamura K, Barzi F, Lam TH, Huxley R, Gu D, Patel A, Kim HC WM. Smoking, diabetes and cardiovascular diseases in men in the Asia Pacific region. *J Diabetes*. 2009;1(3):2009.
80. Huxley R, Barzi F WM. Excess risk of fatal coronary heart disease associated with diabetes in men and women: Meta-analysis of 37 prospective cohort studies. *Br Med Journal*. 2006;332(7533):73–8.
81. Chan JC, Malik V, Jia W, Kadowaki T, Yajnik CS, Yoon KH HF. Diabetes in Asia: Epidemiology, risk factors, and pathophysiology. *J Am Med Assoc*. 2009;301(20):2129–40.
82. Nathan DM, Cleary PA, Backlund JY, Genuth SM, Lachin JM OT. Intensive diabetes treatment and cardiovascular disease in patients with type 1 diabetes. *N Engl J Med*. 2005;353(25):2643–53.
83. Holman RR, Paul SK, Bethel MA, Matthews DR NH. 10-year follow-up of intensive glucose control in type 2 diabetes. *New Engl J Med*. 2008;359(15):1577–89.
84. Kelly TN, Bazzano LA, Fonseca VA, Thethi TK, Reynolds K HJ. Glucose control and cardiovascular disease in type 2 diabetes. *Ann Intern Med*. 2009;151(6):1–10.
85. Arking DE CA. Understanding cardiovascular disease through the lens of genome-wide association studies. *Trends Genet*. 2009;25(9):387–94.
86. Palomaki GE, Melillo S BL. Association between 9p21 genomic markers and heart disease: A meta-analysis. *J Am Med Assoc*. 2010;303(7):648–56.
87. Arnett DK, Baird AE, Barkley RA, Basson CT BE. Relevance of genetics and genomics for prevention and treatment of cardiovascular disease. *Circulation*. 2007;115(22):2878–901.
88. Gluckman PD, Hanson MA, Buklijas T, Low FM BA. Epigenetic mechanisms that underpin metabolic and cardiovascular diseases. *Nat Rev Endocrinol*. 2009;5(7):401–8.
89. G. J. Gender differences in cardiovascular disease prevention. *Menopause Int*. 2008;14(1):213–7.
90. Allen J SS. Gender, ethnicity, and cardiovascular disease. *J Cardiovasc Nursing*. 2005;20(1):1–6.
91. Anand SS, Islam S, Rosengren A FM. Risk factors for myocardial infarction in women and men: Insights from the INTERHEART study. *Eur Hear Journal*. 2008;29(7):932–40.

92. V R-Z. Therapeutic implications of the gender-specific aspects of cardiovascular disease. *Nature*. 2006;5(5):425–38.
93. Buchanan MR BS. Sex-related differences in the pathophysiology of cardiovascular disease: Is there a rationale for sex-related treatments? *Can J Cardiol*. 2001;17((Suppl D)):7D – 13D.
94. Roeters van Lennep JE WH. Risk factors for coronary heart disease: Implications of gender. *Cardiovasc Res*. 2002;53(3):538–49.
95. MJ. L. Cardiovascular disease in women: Gender-specific aspects of hypertension and the consequences of treatment. *J Women’s Heal*. 1999;7(2):199–209.
96. Islam S N, Khan N I AN. Food composition table and database for Bangladesh with special reference to selected ethnic food. Dhaka.: Institute of Nutrition and Food Science, Dhaka University.; 2012.
97. World Health Organization (WHO). Global Physical Activity Questionnaire (GPAQ) Analysis Guide [Internet]. Geneva: World Health Organization. 2012. Available from: www.who.int/chp/steps/resources/GPAQ_Analysis_Guide.pdf
98. National Institute of Alcohol Abuse and Alcoholism. Newsletter. 2004;3.
99. CDC. FAQ Alcohol and Public Health. Available from: <http://www.cdc.gov/alcohol/faqs.htm>
100. Human USD of A and USD of H and S. Dietary Guidelines for Americans. 7th Editio. Washington, DC: US Government Printing Office; 30–32. p.
101. Chobanian AV, Bakris GL, Black HR, Cushman WC, Green LA, Izzo JL Jr et al. Seventh report of the Joint National Committee on Prevention, Detection, Evaluation, and Treatment of High Blood Pressure. *Hypertension*. 42(6):1206–52.
102. Hulzebosch A, van de Vijver S, Oti SO, Egondi T KC. Profile of people with hypertension in Nairobi’s slums: a descriptive study. *Glob Heal*. 2015;11:26.
103. Chaturvedi S, Pant M, Neelam YG. Hypertension in Delhi: Prevalence, awareness, treatment and control. *Trop Doct*. 2007;37:142–5.
104. Anand K, Shah B, Yadav K, Singh R, Mathur P, Paul E et al. Are the Urban poor vulnerable to non-communicable diseases? A survey of risk factors for non-communicable diseases in urban slums of Faridabad. *Natl Med J India*. 2007;20:115–20.

105. Misra A, Pandey RM, Devi JR, Vikram NK KN. High prevalence of diabetes obesity and dyslipidemia in urban slum population of northern India. *Int J Obes Relat Metab Disord.* 2001;11:1722–9.
106. Mohan V, Shanthirani S, Deepa R, Premalatha G, Sastry NG SR. Intra urban differences in the prevalence of the metabolic syndrome in Southern India. The Chennai Urban population Study. (CUPS). *Diabet Med.* 2001;18:280–7.
107. Ongeti KW, Ogengo JA, Pulci AN, Olabu BO GC. Blood pressure characteristics among slum dwellers in Kenya. *Glob Adv Res J Microbiol.* 2013;2:080–5.
108. Joshi MD et al. Prevalence of hypertension and associated cardiovascular risk factors in an urban slum in Nairobi, Kenya: A population-based survey. *BMC Public Health.* 2014;14:1177.
109. Singh R, Mukherjee M, Kumar R, Singh R PR. Study of Risk factors of Coronary Heart Disease in Urban Slums of Patna. *NepalJournal Epidemiol [Internet].* 2012;2(3):205–12. Available from: www.nepjol.info/index.php/NJE
110. Panesar S, Chaturvedi S, Saini N K, Avasthi R SA. Prevalence and predictors of hypertension among residents aged 20-59 years of a slum-resettlement colony in Delhi, India. *WHO South-East Asia J Public Heal.* 2013;2:83–7.
111. Rani M, S Bonu, Jha P, SN Njuyen L. Tobacco use in India: prevalence and predictors of smoking and chewing in a national cross sectional household survey . *Tob Control [online].* 12(4).
112. Yatsuya H FA. Risk of incident cardiovascular disease among users of smokeless tobacco in the Atherosclerosis Risk in Communities (ARIC) study. *Am J Epidemiol.* 2010;172:600–5.
113. Teo KK, Ounpuu S, Hawken S, Pandey MR, Valentin V, Hunt D et al. Tobacco use and risk of myocardial infarction in 52 countries in the INTERHEART study: A case-control study. *Lancet.* 2006;368:647–58.
114. Rahman MA ZM. Smoking and smokeless tobacco consumption: Possible risk factors for coronary heart disease among young patients attending a tertiary care cardiac hospital in Bangladesh. *Public Heal.* 2008;122:1331–8.
115. National cardiovascular disease Database. ICMR Task force project on collaborative study of coronary heart disease.

116. Raju SS K V. Prevalence of Risk Factors for Cardiovascular Disease and Type II Diabetes in Females within the Slum Nuevo Jerusalem, Peru. *Int J Diabetes Clin Res.* 2:028.
117. Colin Bell A, Adair LS PB. Ethnic differences in the association between body mass index and hypertension. *Am J Epidemiol.* 2002;155(4):346–53.
118. Chirinos JA, Franklin SS, Townsend RR RL. Body mass index and hypertension hemodynamic subtypes in the adult US population. *Arch Intern Med.* 2009;169(6):580–6.
119. SJ. G. Through the doors of perception and reality about cardiovascular disease. *Pfizer J.* 2003;7(1).

CHAPTER 8
ANNEXURE

Annex-1

Prevalence of Cardiovascular Disease Risk Factors among Adults in a Slum of Dhaka City							
Work plan (July 2014- June 2015)							
Sl.No.	Activity detail	1-2 months	3-4 months	5-6 months	7-8 months	9-10 months	11-12 months
1	Title selection and approval						
2	Protocol writing						
3	Literature review						
4	Questionnaire Development, Training of the Data collector and Pilot study						
5	Data collection						
6	Data processing and data analysis						
7	Report writing and submission						

Annex-2

Informed Consent Form (English)

Dear Participant

Assalamualaikum, you have been selected to be a part of this thesis and this is why we would like to interview you. This thesis is being conducted by Dr. S. M. Zahid Hassan Arefin, a student of M.Phil in Epidemiology and Biostatistics at Bangladesh Institute of Health Sciences (BIHS) under University of Dhaka. We are interested in conducting a research on Risk factors of Cardiovascular Diseases among adults in your slum. The purpose of this thesis is to find out the prevalence of cardiovascular diseases risk factors among adults in a slum of Dhaka city.

Title of thesis

Prevalence of cardiovascular diseases risk factors among adults in a slum of Dhaka city.

Why you are selected for this study?

This study will assess cardiovascular diseases risk factors among adults in a slum of Dhaka city. You will be one of approximately 221 participants to be asked to participate in this study.

Data Collection Methods

Step 1- Interview questions

Step 2- Physical Measurements

Risks and Discomforts

You will not be charged for any part of the examination. We do not expect an unusual risk to occur as a result of participation. In the unlikely event that during examination procedures if you require medical care, first aid will be available.

Confidentially

The information you provide is totally confidential and will not be disclosed to anyone. It will only be used for research purposes. Your name, address, and other personal information will be removed from the instrument and only a code will be used to connect your name and your answers without identifying you. We will not use your name in sharing and publishing the results of this study.

Voluntary Participation

Your participation is voluntary and you can withdraw from the study after having agreed to participate. You are free to refuse to answer any question that is asked in the questionnaire. If you have any questions about this study you may ask me or contact to the researcher [Dr. Zahid Arefin, cell no: +8801911255960].

Results

The results of this thesis will be used to help plan strategies in reducing the risk factor that contribute to cardiovascular diseases in Bangladesh. The results will be published in research publications, media briefings and reports and can be made available to you on your request.

Consent to Participate

Signing this consent form indicates that you have read this consent form (or have had it read to you) and that you voluntarily agree to participate in this research study.

Signatures

I hereby provide INFORMED CONSENT to take part in Steps 1 and 2 of this Risk Factor Study.

Name of the participant:

Witness:

Sign or fingerprint

Sign or fingerprint

Annex-3
Questionnaire- Bangla

Prevalence of Cardiovascular Disease Risk Factors among Adults in a Slum of Dhaka City

তথ্য প্রদানকারীর আই.ডি.নং	□ □ □ □
সাক্ষাতকারের তারিখ	□ □ □ □ □ □ □ □
তথ্য প্রদানকারীর নাম:	
ঠিকানা:	
ফোন নং (যদি থাকে)	

Step 1 ডেমোগ্রাফিক তথ্যাবলী

মূল: ডেমোগ্রাফিক তথ্যাবলী			
প্রশ্নাবলী		উত্তর	কোড
১	লিঙ্গ	পুরুষ ১ <input type="checkbox"/> মহিলা ২ <input type="checkbox"/>	D1
২	আপনার বয়স কত?	<input type="text"/> বৎসর	D2
৩	সর্বমোট কত বৎসর প্রাতিষ্ঠানিক শিক্ষা গ্রহণ করেছেন? (প্রথম শ্রেণীর নিচে এবং উপানুষ্ঠানিক শিক্ষা অসম্পূর্ণ হলে না)	<input type="text"/> বৎসর	D3

বর্ধিত: ডেমোগ্রাফিক তথ্যাবলী			
প্রশ্নাবলী		উত্তর	কোড
৪	আপনি কতদূর লেখাপড়া করেছেন? কোন প্রাতিষ্ঠানিক শিক্ষা নেই ... ১ প্রাথমিক শিক্ষা অসম্পূর্ণ... ২ প্রাথমিক শিক্ষা... ৩ মাধ্যমিক শিক্ষা... ৪ উচ্চ মাধ্যমিক শিক্ষা... ৫ স্নাতক/ ডিগ্রী... ৬ স্নাতকোত্তর... ৭ জানাতে অসম্মতি ... ৮৮	<input type="text"/>	D4
৫	আপনার ধর্ম কি? ইসলাম ... ১ হিন্দু ... ২ খ্রিষ্টান ... ৩ বৌদ্ধ ... ৪ অন্যান্য ... ৫ জানাতে অসম্মতি ... ৮৮	<input type="text"/>	D5

କ୍ଷମାକରଣ ପରୀକ୍ଷା (Behavioral Measurements)	
କେଉଁ	ଉତ୍ତର
T1	କି? ୧ ଆପଣ କି ବର୍ତ୍ତମାନ କୋଣ ଧ୍ୟାନ କରନ୍ତେ? ସେନ- ନିମ୍ନୋପ, ବିଚି, ଡକ୍ଟା, ପିମ୍ପୁ, ମାହିମ । (ନିୟମା କାଳ ସମାପ୍ତ)
T2	କି? ୧ ଆପଣ କି ଉଚିତ୍ତନ/ନିୟମିତ ଧ୍ୟାନ କରନ୍ତେ?
T3	କି? ୧ କତ ବସନ୍ତ ସମୟ ଥିବେ ଆପଣ ଧ୍ୟାନ କରନ୍ତେ କରନ୍ତେ?
T4a	କି? ୧ କତ ବସନ୍ତ ସମୟ ଧ୍ୟାନ କରନ୍ତେ କରନ୍ତେ?
T4b	କି? ୧ କତ ବସନ୍ତ ସମୟ ଧ୍ୟାନ କରନ୍ତେ କରନ୍ତେ?
T4c	କି? ୧ କତ ବସନ୍ତ ସମୟ ଧ୍ୟାନ କରନ୍ତେ କରନ୍ତେ?
T5a	କି? ୧ ନିମ୍ନୋପ
T5b	କି? ୧ ବିଚି
T5c	କି? ୧ ଡକ୍ଟା/ସାମା
T5d	କି? ୧ ମାହିମ
T5e	କି? ୧ ପିମ୍ପୁ ନିମ୍ନୋପ
T5f	କି? ୧ ପିମ୍ପୁ
T5g	କି? ୧ ଆପଣ
T5	କି? ୧ ଆପଣ (ନିମ୍ନୋପ କରନ୍ତେ)

ଧ୍ୟାନ : ଆପଣଙ୍କର ସାଧାରଣ
 ଏହା ଆପଣଙ୍କୁ ବିଚିତ୍ତନ ସାଧି ବିଷୟକ ଜ୍ଞାନାଗ୍ରହଣ ମନେକରି କିଛି କିଛି କର । ଏହାପରେ ଆପଣଙ୍କୁ ଧ୍ୟାନ, ସମାପନ, କଳ ଓ ନୀଳ-ନୀଳ
 ଧ୍ୟାନ ଏବଂ ଆଧ୍ୟାତ୍ମିକ ମନେକରି କରନ୍ତେ । ଧ୍ୟାନ କରନ୍ତେ ।

ব্যক্তি: তথাকথন ব্যবহার		ব্যক্তি	উত্তর	কোড	
১৬.১	পূর্বে কি আপনাকে এটিদিন/নিয়মিত ব্যবহার করেছিলেন?	হ্যাঁ <input type="checkbox"/> না <input type="checkbox"/>	১ (যদি না হয়, তবে ১৭নং প্রশ্নে যান)	T6	
১৬.২	কতদিন ব্যবহার আপনাকে এটিদিন/নিয়মিত ব্যবহার করেছেন বা? (যে কোন ১টি উত্তর দিন)	কখনো <input type="checkbox"/> ১-৩ দিন <input type="checkbox"/> ৩-৬ দিন <input type="checkbox"/> ৬-১২ দিন <input type="checkbox"/> ১২-২৪ দিন <input type="checkbox"/> ২৪-৬০ দিন <input type="checkbox"/> ৬০-১২০ দিন <input type="checkbox"/> ১২০-১৮০ দিন <input type="checkbox"/> ১৮০ দিন বা তার বেশি <input type="checkbox"/>	কোনো <input type="checkbox"/> ১ (যদি না হয়, তবে ১৭নং প্রশ্নে যান)	T7a T7b T7c	
১৭	আপনি কি বর্তমানে কোন ধর্মীয়/বিশ্বাসী জীবন ত্যাগ করেছেন?	হ্যাঁ <input type="checkbox"/> না <input type="checkbox"/>	১ (যদি না হয়, তবে ২০ নং প্রশ্নে যান)	T9	
১৮	আপনি কি ধর্মীয়/বিশ্বাসী জীবন ত্যাগ করেছেন?	হ্যাঁ <input type="checkbox"/> না <input type="checkbox"/>	১ (যদি না হয়, তবে ২০ নং প্রশ্নে যান)	T10	
১৯	পূর্বে কি এটিদিন কতবার এজেন্সি ব্যবহার করেছেন? (একটিই উত্তর দিনে)	কখনো <input type="checkbox"/> ১-৩ <input type="checkbox"/> ৪-৬ <input type="checkbox"/> ৭-১০ <input type="checkbox"/> ১১-১৫ <input type="checkbox"/> ১৬-২০ <input type="checkbox"/> ২১-৩০ <input type="checkbox"/> ৩১-৪৫ <input type="checkbox"/> ৪৬-৬০ <input type="checkbox"/> ৬০-৭৫ <input type="checkbox"/> ৭৫-৯০ <input type="checkbox"/> ৯১-১০০ <input type="checkbox"/> ১০০-১২০ <input type="checkbox"/> ১২০-১৫০ <input type="checkbox"/> ১৫০-১৮০ <input type="checkbox"/> ১৮০-২০০ <input type="checkbox"/> ২০০-২৫০ <input type="checkbox"/> ২৫০-৩০০ <input type="checkbox"/> ৩০০-৩৫০ <input type="checkbox"/> ৩৫০-৪০০ <input type="checkbox"/> ৪০০-৪৫০ <input type="checkbox"/> ৪৫০-৫০০ <input type="checkbox"/> ৫০০-৫৫০ <input type="checkbox"/> ৫৫০-৬০০ <input type="checkbox"/> ৬০০-৬৫০ <input type="checkbox"/> ৬৫০-৭০০ <input type="checkbox"/> ৭০০-৭৫০ <input type="checkbox"/> ৭৫০-৮০০ <input type="checkbox"/> ৮০০-৮৫০ <input type="checkbox"/> ৮৫০-৯০০ <input type="checkbox"/> ৯০০-৯৫০ <input type="checkbox"/> ৯৫০-১০০০ <input type="checkbox"/>	১ (যদি না হয়, তবে ২০ নং প্রশ্নে যান)	T11a T11b T11c T11d T11e T11f	
২০	আপনাকে এটিদিন কতবার এজেন্সি ব্যবহার করেছেন? (একটিই উত্তর দিনে)	কখনো <input type="checkbox"/> ১-৩ <input type="checkbox"/> ৪-৬ <input type="checkbox"/> ৭-১০ <input type="checkbox"/> ১১-১৫ <input type="checkbox"/> ১৬-২০ <input type="checkbox"/> ২১-৩০ <input type="checkbox"/> ৩১-৪৫ <input type="checkbox"/> ৪৬-৬০ <input type="checkbox"/> ৬০-৭৫ <input type="checkbox"/> ৭৫-৯০ <input type="checkbox"/> ৯১-১০০ <input type="checkbox"/> ১০০-১২০ <input type="checkbox"/> ১২০-১৫০ <input type="checkbox"/> ১৫০-১৮০ <input type="checkbox"/> ১৮০-২০০ <input type="checkbox"/> ২০০-২৫০ <input type="checkbox"/> ২৫০-৩০০ <input type="checkbox"/> ৩০০-৩৫০ <input type="checkbox"/> ৩৫০-৪০০ <input type="checkbox"/> ৪০০-৪৫০ <input type="checkbox"/> ৪৫০-৫০০ <input type="checkbox"/> ৫০০-৫৫০ <input type="checkbox"/> ৫৫০-৬০০ <input type="checkbox"/> ৬০০-৬৫০ <input type="checkbox"/> ৬৫০-৭০০ <input type="checkbox"/> ৭০০-৭৫০ <input type="checkbox"/> ৭৫০-৮০০ <input type="checkbox"/> ৮০০-৮৫০ <input type="checkbox"/> ৮৫০-৯০০ <input type="checkbox"/> ৯০০-৯৫০ <input type="checkbox"/> ৯৫০-১০০০ <input type="checkbox"/>	১ (যদি না হয়, তবে ২০ নং প্রশ্নে যান)	T11a T11b T11c T11d T11e T11f T11g T11h T11i T11j	
২১	আপনাকে এটিদিন কতবার এজেন্সি ব্যবহার করেছেন? (একটিই উত্তর দিনে)	কখনো <input type="checkbox"/> ১-৩ <input type="checkbox"/> ৪-৬ <input type="checkbox"/> ৭-১০ <input type="checkbox"/> ১১-১৫ <input type="checkbox"/> ১৬-২০ <input type="checkbox"/> ২১-৩০ <input type="checkbox"/> ৩১-৪৫ <input type="checkbox"/> ৪৬-৬০ <input type="checkbox"/> ৬০-৭৫ <input type="checkbox"/> ৭৫-৯০ <input type="checkbox"/> ৯১-১০০ <input type="checkbox"/> ১০০-১২০ <input type="checkbox"/> ১২০-১৫০ <input type="checkbox"/> ১৫০-১৮০ <input type="checkbox"/> ১৮০-২০০ <input type="checkbox"/> ২০০-২৫০ <input type="checkbox"/> ২৫০-৩০০ <input type="checkbox"/> ৩০০-৩৫০ <input type="checkbox"/> ৩৫০-৪০০ <input type="checkbox"/> ৪০০-৪৫০ <input type="checkbox"/> ৪৫০-৫০০ <input type="checkbox"/> ৫০০-৫৫০ <input type="checkbox"/> ৫৫০-৬০০ <input type="checkbox"/> ৬০০-৬৫০ <input type="checkbox"/> ৬৫০-৭০০ <input type="checkbox"/> ৭০০-৭৫০ <input type="checkbox"/> ৭৫০-৮০০ <input type="checkbox"/> ৮০০-৮৫০ <input type="checkbox"/> ৮৫০-৯০০ <input type="checkbox"/> ৯০০-৯৫০ <input type="checkbox"/> ৯৫০-১০০০ <input type="checkbox"/>	১ (যদি না হয়, তবে ২০ নং প্রশ্নে যান)	T11j T11k T11l T11m T11n T11o T11p T11q T11r T11s T11t T11u T11v T11w T11x T11y T11z T11aa T11ab T11ac T11ad T11ae T11af T11ag T11ah T11ai T11aj T11ak T11al T11am T11an T11ao T11ap T11aq T11ar T11as T11at T11au T11av T11aw T11ax T11ay T11az T11ba T11bb T11bc T11bd T11be T11bf T11bg T11bh T11bi T11bj T11bk T11bl T11bm T11bn T11bo T11bp T11bq T11br T11bs T11bt T11bu T11bv T11bw T11bx T11by T11bz T11ca T11cb T11cc T11cd T11ce T11cf T11cg T11ch T11ci T11cj T11ck T11cl T11cm T11cn T11co T11cp T11cq T11cr T11cs T11ct T11cu T11cv T11cw T11cx T11cy T11cz T11da T11db T11dc T11dd T11de T11df T11dg T11dh T11di T11dj T11dk T11dl T11dm T11dn T11do T11dp T11dq T11dr T11ds T11dt T11du T11dv T11dw T11dx T11dy T11dz T11ea T11eb T11ec T11ed T11ee T11ef T11eg T11eh T11ei T11ej T11ek T11el T11em T11en T11eo T11ep T11eq T11er T11es T11et T11eu T11ev T11ew T11ex T11ey T11ez T11fa T11fb T11fc T11fd T11fe T11ff T11fg T11fh T11fi T11fj T11fk T11fl T11fm T11fn T11fo T11fp T11fq T11fr T11fs T11ft T11fu T11fv T11fw T11fx T11fy T11fz T11ga T11gb T11gc T11gd T11ge T11gf T11gg T11gh T11gi T11gj T11gk T11gl T11gm T11gn T11go T11gp T11gq T11gr T11gs T11gt T11gu T11gv T11gw T11gx T11gy T11gz T11ha T11hb T11hc T11hd T11he T11hf T11hg T11hh T11hi T11hj T11hk T11hl T11hm T11hn T11ho T11hp T11hq T11hr T11hs T11ht T11hu T11hv T11hw T11hx T11hy T11hz T11ia T11ib T11ic T11id T11ie T11if T11ig T11ih T11ii T11ij T11ik T11il T11im T11in T11io T11ip T11iq T11ir T11is T11it T11iu T11iv T11iw T11ix T11iy T11iz T11ja T11jb T11jc T11jd T11je T11jf T11jg T11jh T11ji T11jj T11jk T11jl T11jm T11jn T11jo T11jp T11jq T11jr T11js T11jt T11ju T11jv T11jw T11jx T11jy T11jz T11ka T11kb T11kc T11kd T11ke T11kf T11kg T11kh T11ki T11kj T11kk T11kl T11km T11kn T11ko T11kp T11kq T11kr T11ks T11kt T11ku T11kv T11kw T11kx T11ky T11kz T11la T11lb T11lc T11ld T11le T11lf T11lg T11lh T11li T11lj T11lk T11ll T11lm T11ln T11lo T11lp T11lq T11lr T11ls T11lt T11lu T11lv T11lw T11lx T11ly T11lz T11ma T11mb T11mc T11md T11me T11mf T11mg T11mh T11mi T11mj T11mk T11ml T11mm T11mn T11mo T11mp T11mq T11mr T11ms T11mt T11mu T11mv T11mw T11mx T11my T11mz T11na T11nb T11nc T11nd T11ne T11nf T11ng T11nh T11ni T11nj T11nk T11nl T11nm T11nn T11no T11np T11nq T11nr T11ns T11nt T11nu T11nv T11nw T11nx T11ny T11nz T11oa T11ob T11oc T11od T11oe T11of T11og T11oh T11oi T11oj T11ok T11ol T11om T11on T11oo T11op T11oq T11or T11os T11ot T11ou T11ov T11ow T11ox T11oy T11oz T11pa T11pb T11pc T11pd T11pe T11pf T11pg T11ph T11pi T11pj T11pk T11pl T11pm T11pn T11po T11pp T11pq T11pr T11ps T11pt T11pu T11pv T11pw T11px T11py T11pz T11qa T11qb T11qc T11qd T11qe T11qf T11qg T11qh T11qi T11qj T11qk T11ql T11qm T11qn T11qo T11qp T11qq T11qr T11qs T11qt T11qu T11qv T11qw T11qx T11qy T11qz T11ra T11rb T11rc T11rd T11re T11rf T11rg T11rh T11ri T11rj T11rk T11rl T11rm T11rn T11ro T11rp T11rq T11rr T11rs T11rt T11ru T11rv T11rw T11rx T11ry T11rz T11sa T11sb T11sc T11sd T11se T11sf T11sg T11sh T11si T11sj T11sk T11sl T11sm T11sn T11so T11sp T11sq T11sr T11ss T11st T11su T11sv T11sw T11sx T11sy T11sz T11ta T11tb T11tc T11td T11te T11tf T11tg T11th T11ti T11tj T11tk T11tl T11tm T11tn T11to T11tp T11tq T11tr T11ts T11tt T11tu T11tv T11tw T11tx T11ty T11tz T11ua T11ub T11uc T11ud T11ue T11uf T11ug T11uh T11ui T11uj T11uk T11ul T11um T11un T11uo T11up T11uq T11ur T11us T11ut T11uu T11uv T11uw T11ux T11uy T11uz T11va T11vb T11vc T11vd T11ve T11vf T11vg T11vh T11vi T11vj T11vk T11vl T11vm T11vn T11vo T11vp T11vq T11vr T11vs T11vt T11vu T11vv T11vw T11vx T11vy T11vz T11wa T11wb T11wc T11wd T11we T11wf T11wg T11wh T11wi T11wj T11wk T11wl T11wm T11wn T11wo T11wp T11wq T11wr T11ws T11wt T11wu T11wv T11ww T11wx T11wy T11wz T11xa T11xb T11xc T11xd T11xe T11xf T11xg T11xh T11xi T11xj T11xk T11xl T11xm T11xn T11xo T11xp T11xq T11xr T11xs T11xt T11xu T11xv T11xw T11xx T11xy T11xz T11ya T11yb T11yc T11yd T11ye T11yf T11yg T11yh T11yi T11yj T11yk T11yl T11ym T11yn T11yo T11yp T11yq T11yr T11ys T11yt T11yu T11yv T11yw T11yx T11yy T11yz T11za T11zb T11zc T11zd T11ze T11zf T11zg T11zh T11zi T11zj T11zk T11zl T11zm T11zn T11zo T11zp T11zq T11zr T11zs T11zt T11zu T11zv T11zw T11zx T11zy T11zz	অন্যান্য <input type="checkbox"/>
অন্যান্য	অন্যান্য <input type="checkbox"/>	হ্যাঁ <input type="checkbox"/> না <input type="checkbox"/>	১ (যদি না হয়, তবে ২০ নং প্রশ্নে যান)	T11	
অন্যান্য	অন্যান্য <input type="checkbox"/>	হ্যাঁ <input type="checkbox"/> না <input type="checkbox"/>	১ (যদি না হয়, তবে ২০ নং প্রশ্নে যান)	T11 others	

		<p>পূর্বে পোষ্যস্বত্বীন তথাহাক ব্যবহার করতেন, কিন্তু বর্তমানে করছেন না এমন কোনো কোনো</p>	
২০	<p>পূর্বে কি কখনো আপন স্বত্বীন তথাহাক ব্যবহার করতেন? (যেমন-ভদ্রা, মাদা-পাতা, জল, নদী)</p>	<p>হ্যাঁ ১ না ২</p>	<input type="checkbox"/>
২১	<p>গত ৭ দিনে, আপন স্বত্বীন তথাহাক ব্যবহার করতেন?</p>	<p>হ্যাঁ ১ না ২ (যদি না হয়, তবে ২০ নং প্রশ্নে যান)</p>	<input type="checkbox"/>
২২	<p>হ্যাঁ হলে গত ৭ দিনের মধ্যে কতদিন আপন স্বত্বীন তথাহাক ব্যবহার করতেন?</p>	<p>দিন</p>	<input type="checkbox"/>
২৩	<p>গত ৭ দিনে, আপন স্বত্বীন তথাহাক ব্যবহার করতেন?</p>	<p>হ্যাঁ ১ না ২ (যদি না হয়, তবে ২০ নং প্রশ্নে যান)</p>	<input type="checkbox"/>
২৪	<p>হ্যাঁ হলে গত ৭ দিনের মধ্যে কতদিন আপন স্বত্বীন তথাহাক ব্যবহার করতেন?</p>	<p>দিন</p>	<input type="checkbox"/>

<p>সূত্র : মদ্যপান</p>		<p>উত্তর</p>
A1a	<p>হ্যাঁ > (যদি না হয়, তবে ১২ নং প্রশ্নে যান)</p>	<p>আপনি কি কখনো মদ্যপান করেছেন?</p> <p>(যেমন- মদ্য, বিয়ার, ওয়াইন, স্পিরিট)</p>
A1b	<p>হ্যাঁ > (যদি না হয়, তবে ১২ নং প্রশ্নে যান)</p>	<p>আপনি কি গত ১২ মাসে কোন ধরনের মদ্যপান করেছেন?</p>
A2	<p>দৈনিক সপ্তাহে ৫-৬ দিন সপ্তাহে ১-৪ দিন মাসে ১-৩ দিন মাসে ১ বা তার কম <input type="checkbox"/></p>	<p>গত ১২ মাসে কোন ধরনের মদ্যপান করেছেন?</p> <p>আপনি কি গত ১২ মাসে কোন ধরনের মদ্যপান করেছেন?</p>
A3	<p>হ্যাঁ > (যদি না হয়, তবে ১২ নং প্রশ্নে যান)</p>	<p>আপনি কি গত ১২ মাসে কোন ধরনের মদ্যপান করেছেন?</p>
A4	<p>হ্যাঁ > (যদি না হয়, তবে ১২ নং প্রশ্নে যান)</p>	<p>গত ১২ মাসে কোন ধরনের মদ্যপান করেছেন?</p>
A5	<p>হ্যাঁ > (যদি না হয়, তবে ১২ নং প্রশ্নে যান)</p>	<p>গত ১২ মাসে কোন ধরনের মদ্যপান করেছেন?</p> <p>এক বার (one single occasion) গুরুত্বপূর্ণ মদ্যপান (standard alcoholic drink=1 unit) মদ্যপান করেছেন?</p> <p>[যদিহ্যাঁ হলে এ প্রশ্নটিতে মদ্যপান করেছেন]</p>
A6	<p>হ্যাঁ > (যদি না হয়, তবে ১২ নং প্রশ্নে যান)</p>	<p>গত ১২ মাসে কোন ধরনের মদ্যপান করেছেন?</p> <p>গত ১২ মাসে কোন ধরনের মদ্যপান করেছেন?</p>

<p>বর্ধিত : খাদ্যাভ্যাস সংকলন তথ্যাবলী</p>		
<p>প্রশ্নাবলী</p>	<p>উত্তর</p>	
<p>৩৬</p> <p>আপনার বাসার বাসার বাস্তব সময় সারাভাগে কোন ধরনের তৈরি ব্যবহার করা হয়? [নামা কাঁচ অথবা] যে কোন একটি নির্দিষ্ট করুন।</p>	<p>১. পরিধান তৈরি..... ২. পায় অংগন..... ৩. পরিষ্কার তৈরি..... ৪. স্যানিটারি তৈরি..... ৫. বাইন বিন তৈরি..... ৬. মাখন/বি..... ৭. ভাজা..... ৮. অন্যান্য..... ৯. নির্দিষ্ট কোনটি নয়..... ১০. কিছই নয়.....</p>	<p>F5</p>
<p>৩৭</p> <p>সারাভাগে আপনার বাসার ১ যানে কি পরিমাণ তৈরি ব্যবহার করা হয়?</p>	<p>কি পরিমাণ</p>	<p>F6</p>
<p>৩৮</p> <p>সারাভাগে আপনার বাসার ১ যানে কি পরিমাণ তৈরি ব্যবহার করা হয়?</p>	<p>কি পরিমাণ</p>	<p>F7</p>
<p>৩৯</p> <p>আপনি কি খাবার সময় সারাভাগে তৈরি তৈরি তৈরি তৈরি তৈরি?</p>	<p>হ্যাঁ ১</p>	<p>F8</p>
<p>৪০</p> <p>বাড়িতে তৈরি হয় না এমন খাবার গড়ে গড়ে কতবার খান? (সকালের নাশা, দুপুর ও রাতের খাবার)</p>	<p>সংখ্যা <input type="text"/> <input type="text"/> জানা না থাকলে ৭৭</p>	<p>F9</p>
<p>Others</p>	<p>.....</p>	

<p>হ্রাস : খাদ্যাভ্যাস সংকলন তথ্যাবলী</p>		
<p>প্রশ্নাবলী</p>	<p>উত্তর</p>	
<p>৩২</p> <p>সারাভাগে সারাভাগে কত দিন ফল খান? [নামা কাঁচ তৈরি তৈরি]</p>	<p><input type="text"/> দিন</p>	<p>F1</p>
<p>৩৩</p> <p>গড়ে একদিনে কতটুকু ফল খান? [নামা কাঁচ অথবা]</p>	<p><input type="text"/> সার্ভিস</p>	<p>F2</p>
<p>৩৪</p> <p>সারাভাগে সারাভাগে কত দিন শাক-সাজি খান? [নামা কাঁচ তৈরি তৈরি]</p>	<p><input type="text"/> দিন</p> <p>জানা না থাকলে ৭৭ (যদি '০০' হয়, তবে ০৬ হবে তৈরি তৈরি)</p>	<p>F3</p>
<p>৩৫</p> <p>গড়ে একদিনে কতটুকু শাক-সাজি খান? [নামা কাঁচ তৈরি তৈরি]</p>	<p><input type="text"/> সার্ভিস</p> <p>জানা না থাকলে ৭৭</p>	<p>F4</p>
<p>খাদ্যাভ্যাস সংকলন তথ্যাবলী</p>	<p>উত্তর</p>	
<p>খাদ্যাভ্যাস সংকলন তথ্যাবলী</p> <p>খাদ্যাভ্যাস সংকলন তথ্যাবলী</p> <p>খাদ্যাভ্যাস সংকলন তথ্যাবলী</p>		

<p>(a-b) P16</p>	<p>যদি <input type="checkbox"/> হ্যাঁ <input type="checkbox"/></p>	<p>১৬.০০ টাকা পর্যন্ত হারে কর প্রদান করা হবে।</p>
<p>১৬.০০ টাকার উপর কর প্রদান করা হবে।</p> <p>১৬.০০ টাকার উপর কর প্রদান করা হবে।</p>		
<p>(a-b) P15</p>	<p>যদি <input type="checkbox"/> হ্যাঁ <input type="checkbox"/></p>	<p>১৫.০০ টাকা পর্যন্ত হারে কর প্রদান করা হবে।</p>
<p>P14</p>	<p>না <input type="checkbox"/></p>	<p>১৪.০০ টাকা পর্যন্ত হারে কর প্রদান করা হবে।</p>
<p>P13</p>	<p>না <input type="checkbox"/> (যদি না হয়, তবে ০২ হারে কর প্রদান করা হবে)</p>	<p>১৩.০০ টাকা পর্যন্ত হারে কর প্রদান করা হবে।</p>
<p>১৩.০০ টাকার উপর কর প্রদান করা হবে।</p>		
<p>(a-b) P12</p>	<p>যদি <input type="checkbox"/> হ্যাঁ <input type="checkbox"/></p>	<p>১২.০০ টাকা পর্যন্ত হারে কর প্রদান করা হবে।</p>
<p>P11</p>	<p>না <input type="checkbox"/></p>	<p>১১.০০ টাকা পর্যন্ত হারে কর প্রদান করা হবে।</p>
<p>P10</p>	<p>না <input type="checkbox"/> (যদি না হয়, তবে ০২ হারে কর প্রদান করা হবে)</p>	<p>১০.০০ টাকা পর্যন্ত হারে কর প্রদান করা হবে।</p>
<p>কোড</p>	<p>০২</p>	<p>১০.০০ টাকার উপর কর প্রদান করা হবে।</p>

<p>ସାଧିତଃ ଭାଷାବୋଧିନି ମହାବିଦ୍ୟାଳୟ ତଥା</p>	
<p>ଆପଣଙ୍କୁ କି ବଡ଼ତାରେ ଭାଷାବୋଧିନିର ସମ୍ପର୍କ ଭାଙ୍ଗିବା ପାଇଁ କେମିତି କୋଳ ଚିକିତ୍ସା ମୋଡ଼େଲ ?</p>	
୩୩	<p>ହଁ > ନା ></p> <p><input type="checkbox"/></p>
୩୩b	<p>ହଁ > ନା ></p> <p><input type="checkbox"/></p>
୩୪	<p>ହଁ > ନା ></p> <p><input type="checkbox"/></p>
୩୫	<p>ହଁ > ନା ></p> <p><input type="checkbox"/></p>

<p>ସାଧିତଃ ଭାଷାବୋଧିନି ମହାବିଦ୍ୟାଳୟ ତଥା</p>	
<p>ଆପଣଙ୍କୁ କି କଥା ଆପଣଙ୍କର କୋଳ ଚିକିତ୍ସା ମୋଡ଼େଲ ?</p>	
୩୧	<p>ହଁ > ନା ></p> <p><input type="checkbox"/></p>
୩୧b	<p>ହଁ > ନା ></p> <p><input type="checkbox"/></p>
୩୨	<p>ହଁ > ନା ></p> <p><input type="checkbox"/></p>

<p>ସାଧିତଃ ଭାଷାବୋଧିନି ମହାବିଦ୍ୟାଳୟ ତଥା</p>	
<p>ଆପଣଙ୍କୁ କି କଥା ଆପଣଙ୍କର କୋଳ ଚିକିତ୍ସା ମୋଡ଼େଲ ?</p>	
୩୧	<p>ହଁ > ନା ></p> <p><input type="checkbox"/></p>
୩୧b	<p>ହଁ > ନା ></p> <p><input type="checkbox"/></p>

<p>ସାଧିତଃ ଭାଷାବୋଧିନି ମହାବିଦ୍ୟାଳୟ ତଥା</p>	
<p>ଆପଣଙ୍କୁ କି କଥା ଆପଣଙ୍କର କୋଳ ଚିକିତ୍ସା ମୋଡ଼େଲ ?</p>	
୩୧	<p>ହଁ > ନା ></p> <p><input type="checkbox"/></p>
୩୧b	<p>ହଁ > ନା ></p> <p><input type="checkbox"/></p>

Questionnaire- English

Prevalence of Cardiovascular Disease Risk Factors among Adults in a Slum of Dhaka City

Participant Identification Number	<input type="text"/> <input type="text"/> <input type="text"/>
Date of interview	<input type="text"/> <input type="text"/> <input type="text"/> <input type="text"/> <input type="text"/> <input type="text"/>
Participant's name	
Address	
Phone No (if available)	

Step 1 Demographic Information

CORE: Demographic Information			
Question	Response	Code	
1	Sex Male 1 <input type="checkbox"/> Female 2 <input type="checkbox"/>	D1	
2.	How old are you? <input type="text"/> <input type="text"/> Years	D2	
3	In total, how many years have you spent at school or in full-time study (excluding less than Class 1 and pre-school)? <input type="text"/> <input type="text"/> Years	D3	

EXPANDED: Demographic Information			
Question	Response	Code	
4	What is the highest level of education you have completed? No formal schooling 1 Less than primary school 2 Primary school completed 3 Secondary school completed 4 Higher Secondary school completed 5 College/University completed 6 Post graduate degree completed 7 Refused 88	<input type="text"/> <input type="text"/>	D4
5	What is your religion ? Islam 1 Hinduism 2 Christianity 3 Buddhism 4 Others 5 Refused 88	<input type="text"/> <input type="text"/>	D5

6	<p>What is your marital status?</p> <p>unmarried 1 married 2 Separated 3 Divorced 4 Widowed 5 Refused 88</p>	<div style="text-align: center;"> <input type="text"/> <input type="text"/> </div>	D6
7	<p>Which of the following best describes your main work status over the past 12 months?</p> <p>Government employee 1 Non-government employee 2 Business (small) 3 Business (large) 4 Farming (Land owner & farmer) 5 Agricultural worker 6 Industrial worker 7 Daily laborer 8 Rickshaw puller 9 Student 10 Housework 11 Retired 12 Unemployed, able to work 13 Unemployed, unable to work 14 Self employed 15 Driver 16 Other 17 Refused to tell 88</p>	<div style="text-align: center;"> <p>498070</p> <input type="text"/> <input type="text"/> </div>	D7
	<p>Please specify, if other</p>	<p>.....</p> <p>..</p>	D7a
8	<p>What is the number of your family member? (exclude children below 6 months of age)</p>	<div style="text-align: center;"> <input type="text"/> <input type="text"/> </div>	D8
9	<p>What is your average monthly family income?</p>	<p>.....</p> <p>Taka.</p>	D9
10	<p>In which type of house do you live in?</p> <p>Kacha (made of clay)..... 1 Tin shade.....2 Brick made.....3 Other.....4</p>	<div style="text-align: center;"> <input type="checkbox"/> </div>	D10
	<p>Please specify, if other</p>	<p>.....</p> <p>..</p>	D10a

Step 1 (Behavioral Measurements)

CORE: Tobacco Use		
Now I am going to ask you some questions about various health behaviours. This includes things like smoking, drinking alcohol, eating fruits and vegetables and physical activity. Let's start with tobacco.		
Question	Response	Code
11 Do you currently smoke any tobacco products , such as cigarettes, cigars or pipes? <i>(as showcard)</i>	Yes 1 <input type="checkbox"/> No 2 <input type="checkbox"/> <i>(If No, go to question 16)</i>	T1
12 Do you smoke tobacco products Daily/ regularly?	Yes 1 <input type="checkbox"/> No 2 <input type="checkbox"/> <i>(If No, go to question 16)</i>	T2
13 How old were you when you first started smoking daily?	Age (years) <input type="text"/> <input type="text"/> Don't know 77	T3
14 Do you remember how long ago it was?	<input type="text"/> <input type="text"/> Years Or Months Or Weeks Don't know 77	T4a T4b T4c
15 On average, how many of the following do you smoke each day?		
Manufactured cigarettes	<input type="text"/> <input type="text"/>	T5a
Biris	<input type="text"/> <input type="text"/>	T5b
Hukkah/Dhaba	<input type="text"/> <input type="text"/>	T5c
Pipes	<input type="text"/> <input type="text"/>	T5d
Hand-rolled cigarettes	<input type="text"/> <input type="text"/>	T5e
Cigars, cheroots	<input type="text"/> <input type="text"/>	T5f
Other	Yes 1 <input type="checkbox"/> No 2 <input type="checkbox"/>	T5g
Please specify, if other	T5 others

EXPANDED: Tobacco Use			
Question	Response	Code	
16.1	In the past, did you ever smoke daily ?	Yes 1 <input type="checkbox"/> No 2 (If No, go to question 17)	T6
16.2	How long ago did you stop smoking daily? (<i>record only 1, not al 3</i>)	Years ago <input type="checkbox"/> <input type="checkbox"/> Months ago Weeks ago Don't know 77	T7a T7b T7c
17	Do you currently use any smokeless tobacco such as zarda, sadapata, gul, snuff?	Yes 1 <input type="checkbox"/> No 2 (If No, go to question 20)	T9
18	Do you currently use smokeless tobacco products such as zarda, sadapata, gul, snuff daily ?	Yes 1 <input type="checkbox"/> No 2 (If No, go to question 20)	T10
19	On average, how many times a day do you use ...		
	zarda	<input type="checkbox"/> <input type="checkbox"/>	T11a
	sadapata	<input type="checkbox"/> <input type="checkbox"/>	T11b
	gul,	<input type="checkbox"/> <input type="checkbox"/>	T11c
	snuff	<input type="checkbox"/> <input type="checkbox"/>	T11d
	Other	Yes 1 <input type="checkbox"/> No 2	T11e
	Other (specify)	T11 others

20	In the past , did you ever use smokeless tobacco such as zarda, sadapata, gul, snuff daily ?	Yes 1 <input type="checkbox"/> No 2 <input type="checkbox"/>	T12
21	During the past 7 days, did someone in your home smoke when you were present?	Yes 1 <input type="checkbox"/> No 2 <input type="checkbox"/> <i>((If No, go to question 23))</i>	T13
22	If yes, on how many days did someone in your home smoke when you were present?	Number of days <input type="checkbox"/>	T14
23	During the past 7 days, did someone smoke in closed areas in your workplace (in the building, in a work area or a specific office) when you were present?	Yes 1 <input type="checkbox"/> No 2 <input type="checkbox"/> <i>((If No, go to question 25))</i>	T15
24	If yes, on how many days did someone smoke in closed areas in your workplace (in the building, in a work area or a specific office) when you were present?	Number of days <input type="checkbox"/>	T16

CORE: Alcohol Consumption			
Question	Response	Code	
25	Have you ever consumed an alcoholic drink such as local wine, beer, wine, spirit?	Yes 1 No 2 (<i>If No, go to question 32</i>) <input type="checkbox"/>	A1a
26	Have you consumed an alcoholic drink within the past 12 months ?	Yes 1 No 2 (<i>If No, go to question 32</i>) <input type="checkbox"/>	A1b
27	During the past 12 months, how frequently have you had at least one alcoholic drink?	Daily 1 5-6 days per week 2 1-4 days per week 3 1-3 days per month 4 Less than once a month 5	A2
28	Have you consumed an alcoholic drink within the past 30 days ?	Yes 1 No 2 (<i>If No, go to question 32</i>) <input type="checkbox"/>	A3
29	During the past 30 days, on how many days did you have at least one alcoholic drink?	<input type="checkbox"/> <input type="checkbox"/> Don't know 77	A4
30	During the past 30 days, what was the average number of standard alcoholic drinks you had (in one single occasion), counting all types of alcoholic drinks together?	<input type="checkbox"/> <input type="checkbox"/> Don't know 77	A5
31	During the past 30 days, how many times did you have for men: five or more, for women: four or more standard alcoholic drinks in a single drinking occasion?	<input type="checkbox"/> <input type="checkbox"/> Don't know 77	A6

CORE: Diet			
The next questions ask about the fruits and vegetables that you usually eat. I have a nutrition card here that shows you some examples of local fruits and vegetables. Each picture represents the size of a serving. As you answer these questions please think of a typical week in the last year.			
Question	Response	Code	
32	In a typical week, on how many days do you eat fruit? (as showcard)	Number of days <input type="text"/> <input type="text"/> Don't Know 77 (If 00, go to question 34)	F1
33	How many servings of fruit do you eat on one of those days? (as showcard)	Number of servings Don't Know 77 <input type="text"/> <input type="text"/>	F 2
34	In a typical week, on how many days do you eat vegetables? (as showcard)	Number of days <input type="text"/> <input type="text"/> Don't Know 77 (If 00, go to question 36)	F 3
35	How many servings of vegetables do you eat on one of those days? (as showcard)	Number of servings Don't know 77 <input type="text"/> <input type="text"/>	F 4

EXPANDED: Diet			
Question	Response	Code	
36	What type of oil or fat is most often used for meal preparation in your household? (USE SHOWCARD) (SELECT ONLY ONE)	Soya bean oil 1 Palm oil 2 Mustard oil 3 Sunflower oil 4 <input type="text"/> <input type="text"/> Rice bran oil 5 Butter or ghee 6 Dalda 7 Other 8 None in particular 9 None used 10	F5
	Other (specify)	F5 Others
37	How much oil is generally used in your home in one month?	<input type="text"/> <input type="text"/> <input type="text"/> <input type="text"/> Liter	F 6
38	How much salt is generally used in your home in one month?	<input type="text"/> <input type="text"/> <input type="text"/> <input type="text"/> Kg	F 7

39	Do you take extra salt while taking food?	Yes 1 No 2	<input type="checkbox"/>	F 8
40	On average, how many meals per week do you eat that were not prepared at home? By meal, I mean breakfast, lunch and dinner.	Number Don't know 77	<input type="checkbox"/> <input type="checkbox"/>	F 9

CORE: Physical Activity

Next I am going to ask you about the time you spend doing different types of physical activity in a typical week. Please answer these questions even if you do not consider yourself to be a physically active person.

Think first about the time you spend doing work. Think of work as the things that you have to do such as paid or unpaid work, study/training, household chores, harvesting food/crops, fishing or hunting for food, seeking employment. In answering the following questions 'vigorous-intensity activities' are activities that require hard physical effort and cause large increases in breathing or heart rate, 'moderate intensity activities' are activities that require moderate physical effort and cause small increases in breathing or heart rate.

Question	Response	Code
41 Does your work involve vigorous-intensity activity that causes large increases in breathing or heart rate like carrying or lifting heavy loads, digging or construction work, harvesting for at least 10 minutes continuously?	Yes 1 No 2 <input type="checkbox"/> <i>(If no, go to question 44)</i>	P1
42 In a typical week, on how many days do you do vigorous-intensity activities as part of your work?	Number of days <input type="checkbox"/>	P2
43 How much time do you spend doing vigorous- intensity activities at work on a typical day?	<input type="text"/> <input type="text"/> <input type="text"/> <input type="text"/> Hour minute	P3 (a-b)
Now I would like to ask you about the time you spend doing moderate-intensity activity		
44 Does your work involve moderate-intensity activity that causes small increases in breathing or heart rate such as brisk walking or carrying light loads for at least 10 minutes continuously? (USE SHOWCARD)	Yes 1 No 2 <input type="checkbox"/> <i>(If no, go to question 47)</i>	P4

45	In a typical week, on how many days do you do moderate-intensity activities as part of your work?	<input type="text"/> days	P5
46	How much time do you spend doing moderate-intensity activities at work on a typical day?	<input type="text"/> <input type="text"/> <input type="text"/> <input type="text"/> Hour minute	P6 (a-b)
Travel to and from places			
47	Do you walk or use a bicycle (pedal cycle) for at least 10 minutes continuously to get to and from places?	Yes 1 No 2 <input type="text"/> (If no, go to question 50)	P7
48	In a typical week, on how many days do you walk or bicycle for at least 10 minutes continuously to get to and from places?	<input type="text"/> days	P8
49	How much time do you spend walking or bicycling for travel on a typical day?	<input type="text"/> <input type="text"/> <input type="text"/> <input type="text"/> hrs mins	P9 (a-b)

CORE: Physical Activity, Continued

Recreational activities

Now I would like to ask you about sports, fitness and recreational activities (leisure)

Question	Response	Code	
50	Do you do any vigorous-intensity sports, fitness or recreational (<i>leisure</i>) activities that cause large increases in breathing or heart rate like running, playing ha-du-du or football for at least 10 minutes continuously? (USE SHOWCARD)	Yes 1 <input type="text"/> No 2 <input type="text"/> (If no, go to question 53)	P10
51	In a typical week, on how many days do you do vigorous-intensity sports, fitness or recreational (<i>leisure</i>) activities?	<input type="text"/> days	P11
52	How much time do you spend doing vigorous-intensity sports, fitness or recreational activities on a typical day?	<input type="text"/> <input type="text"/> <input type="text"/> <input type="text"/> hrs mins	P12 (a-b)

Now I would like to ask you about the time you spend doing moderate-intensity activity outside your work

53	Do you do any moderate-intensity sports, fitness or recreational (<i>leisure</i>) activities that cause a small increase in breathing or heart rate such as brisk walking, cycling, swimming, playing volleyball and dariabandha for at least 10 minutes continuously? (<i>use showcard</i>)	Yes 1 No 2 <input type="checkbox"/> (<i>If no, go to question 56</i>)	P13
54	In a typical week, on how many days do you do moderate-intensity sports, fitness or recreational (<i>leisure</i>) activities?	<input type="text"/> days	P14
55	How much time do you spend doing moderate-intensity sports, fitness or recreational (<i>leisure</i>) activities on a typical day?	<input type="text"/> <input type="text"/> <input type="text"/> <input type="text"/> hrs mins	P15 (a-b)

EXPANDED: Physical Activity

Sedentary behaviour

The following question is about sitting or reclining at work, at home, getting to and from places, or with friends including time spent sitting at a desk, sitting with friends, traveling in car, bus, train, reading, playing cards or watching television, but do not include time spent sleeping.

56	How much time do you usually spend sitting or reclining on a typical day?	<input type="text"/> <input type="text"/> <input type="text"/> <input type="text"/> hrs mins	P16 (a-b)
----	---	---	--------------

CORE: Raised Blood Pressure		
Question	Response	Code
57	Have you ever had your blood pressure measured by a doctor or other health worker? Yes 1 <input type="checkbox"/> No 2 <i>(If no, go to question 61)</i>	H1
58	Have you ever been told by a doctor or other health worker that you have raised blood pressure or hypertension? Yes 1 <input type="checkbox"/> No 2 <i>(If no, go to question 61)</i>	H2

EXPANDED: History of Raised Blood Pressure		
59	Are you currently receiving any treatments for high blood pressure prescribed by a doctor or other health worker? Yes 1 <input type="checkbox"/> No 2	H3a
60	What type of medication are you taking? Alopathic 1 <input type="checkbox"/> Homeo 2 Herbal 3	H3b

CORE: History of Diabetes		
Question	Response	Code
61	Have you ever had your blood sugar measured by a doctor or other health worker? Yes 1 <input type="checkbox"/> No 2 <input type="checkbox"/> <i>(If no, go to question 66)</i>	S1
62	Have you ever been told by a doctor or other health worker that you have raised blood sugar or diabetes? Yes 1 <input type="checkbox"/> No 2 <input type="checkbox"/> <i>(If no, go to question 64)</i>	S2

EXPANDED: History of Diabetes				
Are you currently receiving any of the following treatments/advice for diabetes prescribed by a doctor or other health worker?				
63	Insulin	Yes 1 No 2	<input type="checkbox"/>	S3a
	Oral drugs-modern	Yes 1 No 2	<input type="checkbox"/>	S3b
	Traditional	Yes 1 No 2	<input type="checkbox"/>	S4

Core- Family history of heart attack or stroke				
Question		Response		Code
64	Do you have any family history of heart attack or stroke?	Yes 1 No 2	<input type="checkbox"/>	FH1
65	If yes, specify	heart attack 1 stroke 2	<input type="checkbox"/>	FH 2

Step 2 Physical Measurements

CORE: Height and Weight		
Now I am going to measure your height and weight (in erect posture)		
Question	Response	Code
66	Height In Centimeters (cm) <input style="width: 20px; height: 20px;" type="text"/> <input style="width: 20px; height: 20px;" type="text"/> <input style="width: 20px; height: 20px;" type="text"/> <input style="width: 20px; height: 20px;" type="text"/>	M1
67	Weight In Kilograms (kg) <input style="width: 20px; height: 20px;" type="text"/> <input style="width: 20px; height: 20px;" type="text"/> <input style="width: 20px; height: 20px;" type="text"/> <input style="width: 20px; height: 20px;" type="text"/>	M2
68	For women: Are you pregnant? Yes 1 <input style="width: 20px; height: 20px;" type="checkbox"/> No 2 <input style="width: 20px; height: 20px;" type="checkbox"/> <i>(If no, go to question 70)</i>	M3
CORE: Waist		
Now I am going to measure your waist (in erect posture)		
69	Waist circumference In Centimetres (cm) <input style="width: 20px; height: 20px;" type="text"/> <input style="width: 20px; height: 20px;" type="text"/> <input style="width: 20px; height: 20px;" type="text"/> <input style="width: 20px; height: 20px;" type="text"/>	M4

CORE: Blood Pressure			
Now I am going to measure your blood pressure (in sitting posture)			
70	Reading (After 5 minutes of rest, take the first reading. Let the person take rest again for 5 minutes before taking second reading).	Systolic (mmHg) <input style="width: 20px; height: 20px;" type="text"/> <input style="width: 20px; height: 20px;" type="text"/>	M5a
		Diastolic (mmHg) <input style="width: 20px; height: 20px;" type="text"/> <input style="width: 20px; height: 20px;" type="text"/>	M5b
71	Reading 2 (Take the second reading. If difference between 1 st and 2 nd reading is more than 5 mm Hg, take 3 rd reading. Wait 5 minutes before 3 rd reading).	Systolic (mmHg) <input style="width: 20px; height: 20px;" type="text"/> <input style="width: 20px; height: 20px;" type="text"/>	M6a
		Diastolic (mmHg) <input style="width: 20px; height: 20px;" type="text"/> <input style="width: 20px; height: 20px;" type="text"/>	M6b
72	Reading 3 (Take the third reading)	Systolic (mmHg) <input style="width: 20px; height: 20px;" type="text"/> <input style="width: 20px; height: 20px;" type="text"/>	M7a
		Diastolic (mmHg) <input style="width: 20px; height: 20px;" type="text"/> <input style="width: 20px; height: 20px;" type="text"/>	M7b











Annex-4

SHOWCARDS

A. Tobacco Products

Examples The following picture shows a few selected examples of tobacco products. Sites are to develop show cards including specific examples of local tobacco products.

Step	Section	Items
Step 1, Tobacco use	T	T1 to T14

	Manufactured Cigarettes		Snuff, available in wet and dry form
	Roll-your-own (RYO) cigarettes		Betel nut
	Pipe		Chewing tobacco, e.g., plug, loose-leaf, chimo, toombak, guthha or twist.
	Cigars, e.g., cigarillos, double coronas, cheroots, stumpen, chuttas and dhuntis.		Betel leaf
	Biri		Water pipe, also known as shisha, hookah or hubble-bubble

Annex-4

B. Alcohol Consumption

SHOWCARDS

Step	Section	Items
Step 1, alcohol consumption	A	A1-A7

1 standard drink =



1 Standard bottle
of **regular beer**
(285ml)



1 Single measure
of **spirits** (30ml)



1 Medium size
glass of **wine**
(120ml)



1 Measure of
aperitif (60ml)

Note: Net alcohol content of a standard drink is approximately 10g of ethanol. However, standard drinks in different countries can contain different amounts of ethanol. Therefore, countries may have to adapt this measure according to their own standards and will report it measure if different from the standard mentioned above.

Source: World Health Survey Questionnaire.

Annex-4

C. Diet (Typical Fruit And Vegetables And Serving Sizes)

SHOWCARDS

Step	Section	Items
Step 1, Diet	D	D1 to D6

VEGETABLES



Serving size: One standard serving—80 grams (translated into different units of cups depending on type of vegetable and standard cup measures available in the country).




Vegetables	1 Serving size
Raw green leafy vegetables	1 cup
Other vegetables cooked chopped	½ cup
Vegetable Juice	½ cup

Annex-4

SHOWCARDS

Step	Section	Items
Step 1, Diet	D	D1 to D6

FRUITS



Serving size: One standard serving—80 grams (translated into different units of cups depending on type of vegetable and standard cup measures available in the country).

Fruits	1 Serving size
Apple, Banana, Orange	1 medium size piece
Chopped, Cooked or caned fruit	½ cup
Fruit Juice	½ cup Not artificially flavored